

# PHYSICS 105



Q1) Iodine  $^{131}\text{I}$  is widely used in the treatment and diagnosis of the Thyroid gland. The radius (in fm) of this isotope is:

- A) 131.0
- B) 157.2
- C) 5.2
- D) 5.4
- E) 6.1

Q2) A 55-kg person has absorbed a 20-rad dose. How many joules of energy are deposited in his body?

- A) 1.1
- B) 20
- C) 11
- D) 55
- E) 1100

Q3) The activity of 1 gram of radium  $^{226}_{88}\text{Ra}$  is exactly 1 Ci. The half-life of radium (in years) is:

- A) 226
- B) 1170
- C) 2280
- D) 1580
- E) 1950

Q4) A 70-kg researcher absorbs  $4.5 \times 10^8$  neutrons in a workday, each of energy 1.2 MeV. The relative biological effectiveness (RBE) for these neutrons is 10. What is the equivalent dosage of the radiation exposure for this researcher, in mrem?

- A) 1.2
- B) 0.39
- C) 0.77
- D) 3.7
- E) 12

Q5) A 2.0-mCi source of  $^{32}\text{P}$  is implanted in a tumor to give it a 24-Gy dose. The half-life of  $^{32}\text{P}$  is 14.3 days, and 1mCi delivers 10 mGy/min. How long (in min) should the source remain implanted?

- A) 12
- B) 1200
- C) 2400
- D) 300
- E) 800

Q6) Ionizing radiation can be used on meat products to reduce the levels of microbial pathogens. Assume that for refrigerated meat the upper allowed limit is 3.8 kGy. If a beam of electrons, each of energy 1.6 MeV, irradiates 3.0 kg of beef, how many electrons should the beef mass absorb to reach the upper allowed limit?


- A)  $3.8 \times 10^{10}$
- B)  $4.5 \times 10^{10}$
- C)  $3.8 \times 10^{16}$
- D)  $4.5 \times 10^{16}$
- E)  $1.6 \times 10^{10}$

Q7) A biological tissue of mass  $m$  is exposed to 60 rad of alpha radiation. How many rads of slow neutrons can cause the same damage to the same tissues? (For alpha  $\text{RBE}=20$ , for slow neutrons  $\text{RBE} = 5$ ).

- 1) 240
- B) 300
- C) 60
- D) 360
- E) 1200

Q8) A blood vessel of radius  $r$  splits into two smaller vessels, each of radius  $r/3$ . If the velocity in the larger vessel is  $v$ , then the velocity in each of the smaller vessel is:

- A)  $9v$
- B)  $v/9$
- C)  $2v/9$
- D)  $v$
- E)  $9v/2$



09) Water flows into the top floor of a 16 m high building through a pipe of constant 2 cm diameter. At the base of the building (ground level) the water flows into the pipe at a speed of 60 cm/s where the gauge pressure is 3.2 atm. The gauge pressure (in atm) in the pipe in the top floor is:

- A) 0
- B) 1.65
- C) 2.65
- D) 1.54
- E) 3.2


Q10) The surface of water in a tank supplying water to a house is 7 m above the faucet (حنفية) in the house. If the faucet is 2.0-cm diameter, how long (in s) does it take to fill a 0.25-m<sup>3</sup> container in the house?

- A) 95
- B) 57
- C) 68
- D) 80
- E) 136

Q11) How much force ( $F_M$  in N) must the biceps muscle exert when a 5.0-kg mass is held in the hand with the forearm being in static equilibrium in a horizontal position as in the figure. Assume that the elbow joint, O, is 5 cm far from the point


of application of  $F_M$ , and that the mass of forearm and hand together is 2.0 kg.

- A) 800
- B) 402
- C) 100
- D) 200
- E) 50



Q12) The figure shows a uniform, horizontal beam (length = 8 m, mass = 25 kg) that is pivoted to the wall by a hinge at point O, with its far end supported by a cable that makes an angle of 51° with the horizontal. If a person (mass = 60 kg) stands 3.0 m from the pivot, what is the horizontal component of the hinge force (in N) acting at point O?

- A) 380
- B) 189



- C) 0
- D) 20
- E) 278

Q13) The kinetic energy of a car moving along a horizontal road is 130 kJ. The driver applies the breaks, and the car stops in 20 m. The force of friction (in N) (assumed constant) is:

- A) 260000
- B) 2600
- C) 130000
- D) 6500
- E) 1300

Q14) A 55-kg athlete climbs a 9 m long rope in 10s. His average power output (in W) is

- A) 231
- B) 485
- C) 550
- D) 90
- E) 331

Q15) A 4.0 kg mass is placed on a rough surface that makes an angle of  $20^\circ$  with the horizontal. If the mass is on the verge of motion, then the coefficient of static friction ( $\mu_s$ ) is

- A) 0.36
- B) 0.94
- C) 0.87
- D) 0.11
- E) 0.34

Q16) A student moves 6 m along the positive x-direction, then he turns around and moves 9 m along the negative x-direction. His average velocity (in m/s) over the 7.0 s total interval of motion is:

- A) -3
- B) 0.43

- C) 0.75
- D) 3
- E) -0.43

Q17) A stone is projected vertically upwards with a speed of 12 m/s from the top of an 18 m high building. The time (in s) it takes the stone to reach the ground is:

- A) 4.1
- B) 0.1
- C) 3.5
- D) 3.0
- E) 0.6

ANSWERS:

|       |       |       |       |       |
|-------|-------|-------|-------|-------|
| Q1-E  | Q2-C  | Q3-D  | Q4-A  | Q5-B  |
| Q6-D  | Q7-A  | Q8-E  | Q9-B  | Q10-C |
| Q11-B | Q12-E | Q13-D | Q14-B | Q15-A |
| Q16-E | Q17-C |       |       |       |

Q1) Iodine  $^{131}\text{I}$  is widely used in the treatment and diagnosis of the Thyroid gland.

The radius (in fm) of this isotope is:

- A) 131.0
- B) 157.2
- C) 5.2
- D) 5.4
- E) 6.1

$$A = 131$$

$$r = 1.2 \times 10^{-15} \sqrt[3]{A}$$

$$r = 1.2 \times 10^{-15} \sqrt[3]{131}$$

$$= 6.1 \times 10^{-15} = 6.1 \text{ fm}$$

Q2) A 55-kg person has absorbed a 20-rad dose. How many joules of energy are deposited in his body?

- A) 1.1
- B) 20
- C) 11
- D) 55
- E) 1100

$$m = 55 \text{ kg}$$

$$AD = 20 \text{ rad}$$

$$1 \text{ Gy} = 1 \text{ J/kg} =$$

$$x = 20 \text{ rad}$$

$$x = 0.2 \text{ J/kg}$$

$$1 \text{ kg} \rightarrow 0.2 \text{ J}$$

$$55 \text{ kg} \rightarrow x$$

$$x = 11 \text{ J}$$

# Test Banks - Final - All chapters.

- Q3) The activity of 1 gram of radium  $^{226}_{88}\text{Ra}$  is exactly 1 Ci. The half-life of radium (in years) is:
- A) 226
  - B) 1170
  - C) 2280
  - D) 1580**
  - E) 1950

$$A = \lambda N$$

$$3.7 \times 10^{10} = \lambda \times 0.02661 \times 10^{23}$$

$$\lambda = 139 \times 10^{-13} \text{ s}^{-1}$$

$$T_{1/2} = \frac{\ln(2)}{139 \times 10^{-13}} = 0.004986 \times 10^{13} \text{ s}$$

$$n(\text{Ra}) = \frac{m}{M_r} = \frac{1}{226} = 0.00442 \text{ mol}$$



$$1 \text{ mol} \rightarrow 6.022 \times 10^{23} \text{ atoms (Ra)}$$

$$0.00442 \text{ mol} \rightarrow x$$

$$N = x = 0.02661 \times 10^{23} \text{ atoms (Ra)}$$

$$1 \text{ year} = 60 \times 60 \times 24 \times 365$$

$$1 \text{ year} = 31,536,000 \text{ seconds}$$

$$x = 0.004986 \times 10^{13} \text{ s}$$

$$x = 1580 \text{ years}$$

Q4) A 70-kg researcher absorbs  $4.5 \times 10^8$  neutrons in a workday, each of energy 1.2 MeV. The relative biological effectiveness (RBE) for these neutrons is 10. What is the equivalent dosage of the radiation exposure for this researcher, in mrem?

- A) 1.2**
- B) 0.39
- C) 0.77
- D) 3.7
- E) 12

$$E_{\text{neutrons}} = 1.2 \text{ MeV}$$

$$\text{RBE} = 10$$

$$E_d(\text{neutrons}) = \text{RBE} \times \text{AD}$$

$$= (10)(0.1234 \times 10^{-5})$$

$$= 0.1234 \times 10^{-4} \text{ Sv}$$

$$1 \text{ Sv} = 100 \text{ rem}$$

$$0.1234 \times 10^{-4} \text{ Sv} = x$$

$$x = 0.1234 \times 10^{-2} \text{ rem}$$

$$= 1.2 \text{ mrem}$$

$$\text{AD} = \frac{\text{Energy} \times \text{no. of neutrons}}{\text{mass}}$$

$$= \frac{4.5 \times 10^8 \times 1.2 \times 10^6 \times 1.6 \times 10^{-19}}{70}$$

$$= 0.12342 \times 10^{-5} \text{ J/kg}$$



Q5) A 2.0-mCi source of  $^{32}\text{P}$  is implanted in a tumor to give it a 24-Gy dose. The half-life of  $^{32}\text{P}$  is 14.3 days, and 1mCi delivers 10 mGy/min. How long (in min) should the source remain implanted?

- A) 12
- B) 1200
- C) 2400
- D) 300
- E) 800

$$A = 2 \text{ mCi}, \quad 1 \text{ mCi} \rightarrow 10 \text{ mGy/min}$$

$$AD = 24 \text{ Gy}, \quad t_{1/2} = 14.3 \text{ days}$$

$$1 \text{ mCi} \rightarrow 10 \text{ mGy/min}$$

$$2 \text{ mCi} \rightarrow x$$

$$\text{rate} = \frac{\text{dose}}{\text{time}}$$

$$x = 20 \text{ mGy/min}$$

$$= \frac{20 \times 10^{-3} \text{ Gy}}{1 \text{ min}}$$

$$\text{time} = \frac{24}{20 \times 10^{-3}} = 1.2 \times 10^3 = 1200 \text{ min}$$

Q6) Ionizing radiation can be used on meat products to reduce the levels of microbial pathogens. Assume that for refrigerated meat the upper allowed limit is 3.8 kGy. If a beam of electrons, each of energy 1.6 MeV, irradiates 3.0 kg of beef, how many electrons should the beef mass absorb to reach the upper allowed limit?

- A)  $3.8 \times 10^{10}$
- B)  $4.5 \times 10^{10}$
- C)  $3.8 \times 10^{16}$
- D)  $4.5 \times 10^{16}$
- E)  $1.6 \times 10^{10}$

$$E_{\text{electron}} = 1.6 \text{ MeV}, \quad m = 3 \text{ kg beef}$$

(1 eV)

$n \rightarrow$  no. of electrons absorbed by the beef

$$AD_{\text{(max)}} = 3.8 \times 10^3 \text{ Gy}$$

$$\rightarrow AD = \frac{\text{Energy per electron} \times \text{no. of electrons}}{\text{mass}}$$

$$3.8 \times 10^3 = \frac{1.6 \times 10^6 \times 1.6 \times 10^{-19} \times n}{3}$$

$$n = 4.5 \times 10^{16} \text{ electrons}$$

Q7) A biological tissue of mass  $m$  is exposed to 60 rad of alpha radiation. How many rads of slow neutrons can cause the same damage to the same tissues? (For alpha RBE=20, for slow neutrons RBE = 5).

- 1) 240
- B) 300
- C) 60
- D) 360
- E) 1200

$AD = 60 \text{ rad of } \alpha\text{-radiation}$

$RBE(\alpha) = 20 \quad , \quad RBE(\text{neutrons}) = 5$


$ED(\alpha) = ED(\text{neutrons})$

$(20)(60) = (5)(AD)$

$AD = 240 \text{ rad}$

Q8) A blood vessel of radius  $r$  splits into two smaller vessels, each of radius  $r/3$ . If the velocity in the larger vessel is  $v$ , then the velocity in each of the smaller vessel is:

- A)  $9v$
- B)  $v/9$
- C)  $2v/9$
- D)  $v$
- E)  $9v/2$




$A_1 v_1 = A_2 v_2$

$\cancel{\pi} \times (r)^2 v_1 = \cancel{\pi} \left(\frac{r}{3}\right)^2 \times v$

$v_1 = \frac{v}{9}$

Q9) Water flows into the top floor of a 16 m high building through a pipe of constant 2 cm diameter. At the base of the building (ground level) the water flows into the pipe at a speed of 60 cm/s where the gauge pressure is 3.2 atm. The gauge pressure (in atm) in the pipe in the top floor is:

- A) 0
- B) 1.65
- C) 2.65
- D) 1.54
- E) 3.2



$P_1 + \cancel{\rho g h_1} + \frac{1}{2} \cancel{\rho v_1^2} = P_2 + \rho g h_2 + \frac{1}{2} \cancel{\rho v_2^2}$

$P_1 = P_2 + \rho g h_2$

$324160 = P_2 + 156800$

$P_2 = 167360 \text{ Pa}$


$= 1.65 \text{ atm}$

skipl:-  $A_1 v_1 = A_2 v_2$

$A$  is constant, so  $v$  is constant

Q10) The surface of water in a tank supplying water to a house is 7 m above the faucet (حنفية) in the house. If the faucet is 2.0-cm diameter, how long (in s) does it take to fill a 0.25-m<sup>3</sup> container in the house?

- A) 95
- B) 57
- C) 68
- D) 80
- E) 136



$$v_2 = \sqrt{2gh} = \sqrt{2 \times 9.8 \times 7} = 11.7 \text{ m/s}$$


$$\frac{\Delta V}{\Delta t} = A_2 v_2$$

$$\hookrightarrow \frac{0.25}{\Delta t} = \pi \times (1 \times 10^{-2})^2 \times 11.7$$

$$\Delta t = 68 \text{ s}$$

Q11) How much force ( $F_M$  in N) must the biceps muscle exert when a 5.0-kg mass is held in the hand with the forearm being in static equilibrium in a horizontal position as in the figure. Assume that the elbow joint, O, is 5 cm far from the point of application of  $F_M$ , and that the mass of forearm and hand together is 2.0 kg.

- A) 800
- B) 402
- C) 100
- D) 200
- E) 50




$$F_M (5 \times 10^{-2}) = (2 \times 9.8 \times 15 \times 10^{-2}) + (5 \times 9.8 \times 35 \times 10^{-2})$$

$$F_M = 402 \text{ N}$$

Q12) The figure shows a uniform, horizontal beam (length = 8 m, mass = 25 kg) that is pivoted to the wall by a hinge at point O, with its far end supported by a cable that makes an angle of  $51^\circ$  with the horizontal. If a person (mass = 60 kg) stands 3.0 m from the pivot, what is the horizontal component of the hinge force (in N) acting at point O?

- A) 380  
B) 189  
E) 278



$$(60g)(3) + (25g)(4) = (T \sin 51)(8)$$

$$T = 441.35 \text{ N}$$

$$\text{Horizontal component} = T \cos 51^\circ = 278 \text{ N}$$

Q13) The kinetic energy of a car moving along a horizontal road is 130 kJ. The driver applies the breaks, and the car stops in 20 m. The force of friction (in N) (assumed constant) is:

- A) 260000  
B) 2600  
C) 130000  
D) 6500  
E) 1300

$$\cdot KE = 130 \text{ kJ}$$

$$\cdot \text{stopping distance} = 20 \text{ m}$$

$$\cdot f = ?$$

$$W_{nc} = \Delta K$$

$$f \cdot (20)(-1) = 130 \times 10^3$$

$$f = -6.5 \times 10^3 = -6500 \text{ N}$$

Q14) A 55-kg athlete climbs a 9 m long rope in 10s. His average power output (in W) is

- A) 231  
B) 485  
C) 550  
D) 90  
E) 221


$$m = 55 \text{ kg}, d = 9 \text{ m}, \Delta t = 10 \text{ s}$$

$$P = \frac{W}{t} = \frac{Fg \cdot d \cdot \cos 90^\circ}{t} = \frac{(55)(9.8)(9)}{10} = 485 \text{ W}$$

Q15) A 4.0 kg mass is placed on a rough surface that makes an angle of  $20^\circ$  with the horizontal. If the mass is on the verge of motion, then the coefficient of static friction ( $\mu_s$ ) is

- A) 0.36
- B) 0.94
- C) 0.87
- D) 0.11
- E) 0.34

$$m = 4 \text{ kg}$$




$$F_g \sin 20 + f_{\max} = 0$$

$$(\cancel{4})(\cancel{9.8})(\sin 20) = (\mu_s)(\cancel{4})(\cancel{9.8})(\cos 20)$$

$$\mu_s = 0.36$$

Q16) A student moves 6 m along the positive x-direction, then he turns around and moves 9 m along the negative x-direction. His average velocity (in m/s) over the 7.0 s total interval of motion is:

- A) -3
- B) 0.43



$$\bar{v} = \frac{-3}{7} = -0.43$$

1) The engine of a truck of mass 940 kg can deliver an average power of 104800 W. If the truck accelerates from rest, the speed (in m/s) after 4.5 s is: (Ignore air resistance)

- A) 31.7
- B) 36.6
- C) 4.8
- D) 11.2
- E) 15.1

Answer: A

2) A 1 kg ball is located at the top of a 4 m plane inclined at  $45^\circ$  as shown. The ball begins to slide down the inclined plane from rest. The upper half of the inclined plane is frictionless, while the lower half is rough, with a coefficient of kinetic friction  $\mu_k = 0.3$ . The speed (in m/s) of the ball at the bottom of the inclined plane is:

- A) 1.1
- B) 6.9
- C) 7.5
- D) 0.3
- E) 5.3



Answer: B

3) A box of mass  $m$  at a height  $h$  above the floor has a speed  $v$ . Its total mechanical energy is  $E$ . A second box of mass  $m$  at a height  $4h$  above the floor has a speed  $2v$ . The total mechanical energy for the second box is:

- A)  $E$
- B)  $4E$
- C)  $(2)^{1/2}E$
- D)  $(2)^{-1/2}E$
- E)  $2E$

Answer: B

4) A horse drags a heavy cart (200 kg) horizontally on a rough floor at a constant speed. The power delivered by the horse is 1.06 hp. The coefficient of kinetic friction between the cart and the floor is 0.115. The speed (in m/s) with which the cart moves across the floor is: Hint:  $1 \text{ hp} = 746 \text{ W}$

- A) 11.7
- B) 3.5
- C) 2.1
- D) 0.3
- E) 9.0

Answer: B

- 5) When a ball rises vertically to a height  $3h$  and returns to its original position, the work done on it by the gravitational force is :
- A)  $+6mgh$
  - B)  $-3mgh$
  - C)  $+3mgh$
  - D)  $-6mgh$
  - E) Zero

Answer: E


- 6) A motor lifts a 3000 kg elevator 210 m up during a time interval  $t$  at constant speed. If the rate at which the motor does work on the elevator is 362 hp, the time interval  $t$  (in s) is: Hint: 1 hp = 746 w
- A) 1.7
  - B) 23
  - C) 19.9
  - D) 5
  - E) 14.8

Answer: B

- 7) The figure shows a PHY 105 student with a mass of 83 kg. Determine the change in the total mechanical energy (in 104 J) between the initial state (speed of 6.5 m/s and height of 50 m) and the final state (speed of 4.5 m/s and height of 28 m).

- A) +1.53
- B) -1.89
- C) -2.25
- D) -2.36

Answer: B



- 8) A box of mass 18 kg is dropped from rest from a height of 80 m above the floor. The box falls vertically downward and reaches the floor with a speed of 15 m/s. The work (in 103 J) exerted by the air resistance force on the box is:

- A) -12
- B) -14
- C) +12
- D) -16
- E) +16

Answer: A

- 9) A 0.5 kg ball thrown vertically upward with an initial speed of 4.00 m/s has reached a maximum height of 0.8 m. What change does air resistance cause in the mechanical energy (in J) of the ball during the upward motion?
- A) 0.08
  - B) 3.92
  - C) 16
  - D) 4.9

Answer: A

- 10) As shown, a child whose weight is 267 N moves down a distance  $d = 6.1$  m along a slide that makes an angle of  $20.00^\circ$  with the horizontal. If the coefficient of kinetic friction between the slide and the child is 0.1, the change in kinetic energy (in J) of the child over the distance  $d$  is approximately:

- A) 404
- B) 659
- C) 222
- D) 710



Answer: A


- 11) A horse drags a heavy cart (200 kg) horizontally on a rough floor at a constant speed. The power delivered by the horse is 1.06 hp. The coefficient of kinetic friction between the cart and the floor is 0.115. The speed (in m/s) with which the cart moves across the floor is: Hint: 1 hp = 746 W

- A) 3.5
- B) 0.3
- C) 11.7
- D) 2.1
- E) 9.0

Answer: A

- 12) There are two forces acting upon a box as it moves down an incline from point A to point B 2 N applied force directed down the incline and 10 N frictional force. Points A and B are 5 m apart, If the kinetic energy of the box increases by 35 J between A and B, the change in the gravitational potential energy (in J) between to B is:

- A) -75
- B) +75
- C) -10
- D) +95
- E) -95



Answer: B



13) Take car 1 and car 2 car 1 has twice the mass of car 2 but only half the kinetic energy Of car 2. When both cars increase their speed by 5,00 m/s, they then have the same kinetic energy. Calculate the original speed (in m/s) of the car 2.

- A) 7.07
- B) 5.00
- C) 3.53
- D) 11.02
- E) 22.04


Answer: C

14) A 3 kg ball thrown vertically upward has reached a height of 100 m in the presence of air resistance. The air resistance has performed -800 J of work on the ball. Determine the height (in m) the ball would reach if air resistance can be neglected.

- A) 163
- B) 127
- C) 100
- D) 201
- E) 196

Answer: B


15) As shown, a bead of mass 0.5 kg immersed in a certain liquid is released from rest at point A. At point B, the bead has a speed of 6 m/s. The work done on the bead (in J) by the viscosity (friction force) of the liquid is:



- A) +9
- B) -15
- C) -5.7
- D) -9

Answer: C

16) As shown, a horizontal force F is pushing a 1.4 kg block up a frictionless  $14^\circ$  incline from point A to point B which are 1.2 m apart. The work exerted by F on the block is 5 J. If the kinetic energy at point B is 4 J, the kinetic energy (in J) at point A is :




- A) 0
- B) 4
- C) 7.2
- D) 3
- E) 5

Answer: D

- 17) As shown, 2 kg block slides along the track with an initial speed  $V_0$  of 6 m/s. The blue section of the track is frictionless, while the horizontal brown section is rough. On the rough section, a frictional force stops the block in a distance  $d$ . If the height difference  $h$  is 1.1 m and is a coefficient of kinetic friction 0.60, what is  $d$  (in m)?

- A) 4.5
- B) 3.4
- C) 1.2
- D) 5.7
- E) 2.6


Answer: C



- 18) The cart shown is heading left towards a wall, colliding with it and bouncing back to the right. The loss in the mechanical energy (in J) during the bounce is: (Assume that right is the positive direction in the coordinate system.)

- A) 15
- B) 20
- C) 5
- D) 30
- E) 25

Answer: A




- 19) As shown a 2-kg block slides from rest down a frictionless hill at height  $d = 40$  cm. It then moves along a horizontal rough level of length  $d$ , where the coefficient of kinetic friction is 0.50. If the block is still moving it then slides down a second frictionless hill at height  $d/2$  and onto a second horizontal rough level which has length  $d/2$  and where the coefficient of kinetic friction is again 0.50. If the block is still moving, it then slides up a frictionless incline until it momentarily stops. If the block can reach the incline, what is its maximum height (in cm) on the incline measured from the second horizontal level?

The kinetic energy of the block is dissipated entirely into thermal energy along the second horizontal rough level before starting to slide up the incline.

- A) 10
- B) 30
- C) 20
- D) 50


Answer: C



- 20) The figure Shows constant force  $F_0$  (82.0 N) acts on a box (3.00 kg) at angle  $53.00^\circ$ . AS a result, the box moves up the frictionless hill at a constant speed. The work (In J) exerted by  $F_0$  on the box when the box has inclined a vertical distance  $h = 0.150$  m is:

- A) 4.41
- B) 9.8
- C) 7.4
- D) ZERO
- E) 12.3


Answer: A



- 21) The assembly shown is in a complete static equilibrium. Blocks A and B weigh 40 N and 50 N respectively. The string  $T_2$  is exactly horizontal. while the angle  $= 35^\circ$ , The tension  $T_3$  (in N) is:

- A) 2.9
- B) 57.3
- C) 13.3
- D) 50.0
- E) 48.4


Answer: B



- 22) As shown, three forces of equal magnitude act on an object at the origin, The force, points into the screen, Rank the magnitudes of the torques created by these forces at point P1 in descending order (largest first):

- A)  $F_3, F_2, F_1$
- B)  $F_2, F_1, F_3$
- C)  $F_3 = F_2 = F_1$
- D)  $F_1, F_3, F_2$


Answer: B



- 23) A 125 kg cart initially at rest is pulled by three ropes as shown, When the cart moves 100 m horizontally on a frictionless level, it's final speed (in m/s) is:

- A) 19
- B) 27
- C) 22
- D) 30
- E) 24


Answer: E



- 24) The figure shows two blocks released from rest: block A (1.0 kg) and block B (2.0 kg). The frictionless surface inclined at angle  $30^\circ$ . If the pulley has negligible mass, what is the total kinetic energy of the two blocks (in J) when block B has fallen 25 cm?

- A) 0.5
- B) 2.78
- C) 3.68
- D) 7.35
- E) 6.13


Answer: D



- 25) The speed of the truck just before it goes up a frictionless hill (angle:  $15^\circ$ ) is 130 km/h. The truck's mass is  $1.2 \times 10^4$  kg. The minimum length of the hill,  $L$  (in m), needed so that the truck will momentarily stop is:

- A) 1800.8
- B) 1048.7
- C) 256.8
- D) 66.5
- E) 13.3

Answer: C



**Information:**

$R = 8.314 \text{ J/mole.K}$ ;  $k_B = 1.38 \times 10^{-23} \text{ J/K}$ ;  $g = 9.8 \text{ m/s}^2$ .  $\rho_{\text{water}} = 1000.0 \text{ kg/m}^3$  and  $P_{\text{atm}} = 1.013 \times 10^5 \text{ Pa}$ .  
 $1u = 1.66 \times 10^{-27} \text{ kg}$ .  $N_A = 6.02 \times 10^{23} \text{ molecules/mole}$ . Note: Some Results Are Rounded.

1) A patient is administered ( $^{131}\text{I}$ ). How long will it take for the observed radioactivity in her body to decrease to one-fourth its original magnitude? Given that ( $^{131}\text{I}$ ) has half-life ( $T_{1/2}$ ) of 8.1 days

- A) 8.1 days      B) 360 days      C) 376.2 days      **D) 16.2 days**      E) 7.75 days

3) A submarine deep below the surface of the sea is at a gauge pressure of 40 atm. The air inside the submarine is at normal atmospheric pressure. The *net* force (in N) on a flat hull plate 2m by 6m is:

- A)  $4.86 \times 10^2$       B) 4.86      **C)  $4.86 \times 10^7$**       D) 4.92      E)  $4.92 \times 10^7$

4) The linear expansion coefficient for Al is  $\alpha = 2.2 \times 10^{-5} \text{ K}^{-1}$ . What is the increase in volume of a block of  $1 \text{ m}^3$  of Al if the temperature of the block is raised by  $10 \text{ }^\circ\text{C}$ ?

- A)  $220 \text{ cm}^3$       B)  $440 \text{ cm}^3$       C)  $22 \text{ cm}^3$       **D)  $660 \text{ cm}^3$**       E)  $66 \text{ cm}^3$

5) What volume fraction of a cube of density ( $\rho = 0.50 \text{ g/cm}^3$ ) would sink under the surface of a liquid of density ( $\rho_o = 1.0 \text{ g/cm}^3$ )?

- A) 0.80      B) 0.67      C) 0.33      **D) 0.50**      E) 0.20

6) A 63-kg researcher absorbs  $2.6 \times 10^8$  neutrons in a work day. The energy of each neutron is 6.5 MeV. The quality factor (QF) for fast neutrons is 10. The biologically equivalent dosage of the radiation, in mrem (mrem =  $10^{-3} \text{ rem}$ ), is closest to (Note:  $1 \text{ rad} = 0.01 \text{ J/kg}$  and  $1 \text{ ev} = 1.6 \times 10^{-19} \text{ J}$ )


- A) 43      B) 1.3      C) 2.7      D) 13      **E) 4.3**

7) A man pulls a box weighting 40 N a distance of 10 m across the floor at constant speed. How much work (in J) does he do if the coefficient of kinetic friction is 0.20?

- A) 80**      B) -40      C) 0.0      D) 40      E) -80

8) The gravitational force exerted on a solid object, in air, is 4.0 N (Figure a). When the object is suspended from a spring scale and submerged in water, the scale reads 2.0 N (Figure b). Find the density of the object (in  $\text{kg/m}^3$ ). Assume density of water  $\rho = 1000.0 \text{ kg/m}^3$ .

- A) 4000      **B) 2000**      C) 5000  
D) 1000      E) 1500



9) When a man stands, his brain is 0.5 m above his heart. If he bends so that his brain is 0.4 m below his heart, by how much does the blood pressure in his brain change? (Assume density of blood is  $1059.5 \text{ kg/m}^3$ .)

- A) 13.3 kPa    B) 4.0 kPa    C) 13.1 kPa    **D) 9.3 kPa**    E) 16.6 kPa
- 

10) If both gases  $\text{H}_2$  and  $\text{CO}_2$  are at the same temperature. Then the ratio of the *rms* velocities of  $\text{H}_2$  and  $\text{CO}_2$ , [ $V_{\text{rms}}(\text{H}_2)/V_{\text{rms}}(\text{CO}_2)$ ] is: (Given that the molecular mass of  $\text{H}_2 = 2.016 \text{ u}$  and for  $\text{CO}_2 = 44.009 \text{ u}$ )

- A) 21.8    B) 0.21    C) 4.0    D) 0.05    **E) 4.67**
- 

11) Water flows (streamline, nonviscous) from point *a* to point *b* in the horizontal section shown in the figure. Which of the following statements is correct regarding the velocity *v*, pressure *P*, and flow rate *Q* at the two ends of the section?

- A)  $v_a < v_b$ .    B)  $P_a > P_b$     **C)  $P_a < P_b$ .**  
D)  $P_a = P_b$ .    E)  $Q_a > Q_b$  (*Q* is the flow rate).



12)  $^{60}\text{Co}$  beta decays with half life of 5.27 years ( $1.66 \times 10^8 \text{ sec}$ ) into  $^{60}\text{Ni}$ , which then promptly emits gamma rays. These gamma rays are widely used in treating cancer. What is the mass (in gram) of a 1000-Ci cobalt source? (Given that one mole of  $^{60}\text{Co}$  has a mass of 60 g)

- A) 0.118    B) 0.441    **C) 0.882**  
D) 0.245    E) 0.0147
- 

13) If an object was thrown vertically from the ground level with initial speed 25 m/s and return to the same ground level after 5.1 seconds. What is the average velocity (in m/s) of the object when reaching the ground?

- A) 12    B) 24    C) 6    **D) 0**    E) -12
- 

14) The maximum permissible workday dose for occupational exposure to radiation is 26 mrem. A 63 kg laboratory technician absorbs 2.1 mJ of 0.7 MeV gamma rays in a work day. The quality factor (QF) for gamma rays is 1.0. The ratio of the equivalent dosage received by the technician to the maximum permissible equivalent dosage is closest to: (mrem =  $10^{-3}$  rem, 1rad = 0.01 J/kg and 1ev =  $1.6 \times 10^{-19}$  J)


- A) 0.18    B) 0.14    C) 0.17    **D) 0.13**    E) 0.15
- 

15) A radioactive source emits 2.4 MeV neutrons at the rate of 9200 neutrons per second. The number of atoms in the source is  $4.0 \times 10^9$ . The activity of the source, in nCi, is closest to: Hint (nCi =  $10^{-9}$  Ci) and (1Ci =  $3.70 \times 10^{10}$  decays/sec.)

- A) 2500    B) 92    C) 920    D) 25    **E) 250**
-

- 16) The level of the fluid with density  $\rho_s = 1000 \text{ kg/m}^3$  in the left arm of the manometer is 0.2 m above the manometer fluid of density  $\rho_f = 800 \text{ kg/m}^3$  in the right arm. Which of the following relations is true? (Use ;  $g = 10 \text{ m/s}^2$ )

- A.  $P_G = P_{atm}$ .  
 B.  $P_G$  is 2000 Pa higher than  $P_{atm}$ .  
 C.  $P_G$  is 2000 Pa lower than  $P_{atm}$ .  
 D.  $P_G$  is 4000 Pa higher than  $P_{atm}$ .  
 E.  $P_G$  is 6000 Pa higher than  $P_{atm}$



- 17) The radioactive nuclide  $^{60}\text{Co}$  is widely used in medical applications. It undergoes beta decay, and the energy of the decay process is 2.82 MeV per decay event. The half-life of this nucleus is 272 days. Suppose that a patient is given a dose of 6.9 microCurie of  $^{60}\text{Co}$ . If all of this material decayed while in the patient's body, what would be the total energy (in J) deposited there? Hint: ( $1\text{Ci} = 3.70 \times 10^{10}$  decays/sec.) and  $1\text{eV} = 1.6 \times 10^{-19}$  J.

- A) 3.9      B) 11.0      C) 14.0      D)  $8.63 \times 10^{12}$       E)  $4.15 \times 10^6$

- 18) A collapsible plastic bag contains glucose. If the average gauge pressure in the vein is  $1.33 \times 10^3$  Pa, what must be the minimum height  $h$  (in m) of the bag in order to infuse glucose into the vein? Assume density of the solution is equal  $1.02 \rho_{\text{water}}$ .

- A) 0.133      B) 0.113      C) 0.150      D) 0.752      E) 0.333



19. Oxygenation of the deep waters in a sea occurs in early winter due to:

- a. Diffusion of air molecules through water.  
 b. Water mixing resulting from the decrease in density of water at lower as the temp decreases.  
 c. Water mixing resulting from the increase in density of water at lower as the temp decreases.  
 d. The lower density of ice relative to water.  
 e. Water mixing resulting from turbulence and the see waves in early winter.

20. One mole of an ideal gas has a temperature of  $25^{\circ}\text{C}$ . If the volume is held constant and the pressure is doubled, the final temperature (in  $^{\circ}\text{C}$ ) will be

- a. 174      b. 323      c. 50      d. 596      e. 25

21. If water is to be pumped into a water tank at the top of a 10 m high building, what should the water pressure at the base of the building be if the speed of water is constant through the water pipe? ( $1.013 \text{ bar} = 1 \text{ atm}$ ,  $g = 9.8 \text{ m/s}^2$ )


- a. 1.0 bars      b. 2.0 bars      c. 0.5 bars      d. 3.0 bars      e. 0.3 bars

22. The temperature of an object is  $80^{\circ}\text{F}$ . What is its absolute temperature on the Kelvin scale?

- a. 300 K.      b. 335 K.      c. 359 K.      d. 475 K.      e. 400 K.


23. A uniform 100 N beam is held in a vertical position by a pin ( $P$ ) at its lower end and a cable at its upper end. A horizontal force of magnitude  $F = 75 \text{ N}$  acts as shown in the figure. What is the tension in the cable?

- a. 47 N      b. 69 N  
c. 61 N      d. 94 N  
e. 54 N




24. If two objects  $M_1, M_2$  ( $M_1 = M_2$ ) are connected by a light inextensible cord which is attached to the ceiling of an elevator that is accelerating upward at  $2 \text{ m/s}^2$ , the ratio  $T/N$

- a.  $5/3$       b. 2      c. 1      d.  $3/2$       e.  $1/2$




25. The frictional force between mass  $2M$  and the surface is zero, and the frictional force between masses  $M$  and  $2M$  causes both masses to move together when the  $F = 1.2 \text{ N}$  is applied to  $2M$ . If  $M = 1 \text{ kg}$ , what is the frictional force exerted by the large block on the small block?

- a. 0.4 N to the left      b. 0.8 N to the right  
c. 0.4 N to the right      d. 0.8 to the left  
e. 1.2 to the right



26. A block slides on a rough horizontal surface from point A to point B. A force ( $P = 2.0 \text{ N}$ ) acts on the block between A and B, as shown. Points A and B are 1.5 m apart. If the kinetic energies of the block at A and B are 5.0 J and 4.0 J, respectively, how much work is done on the block by the force of friction as the block moves from A to B?


- a.  $-3.3 \text{ J}$       b.  $+1.3 \text{ J}$       c.  $+3.3 \text{ J}$   
d.  $-1.3 \text{ J}$       e.  $+4.6 \text{ J}$





27) Consider the model shown in Figure (b) for a person bending forward to lift a 100-N object. The spine and upper body are represented as a uniform horizontal rod of weight 400 N and length  $L$ , pivoted at the base of the spine. The erector spinal muscle, attached at a point  $2L/3$  away from the pivot, maintains the position of the back. The angle between the spine and this muscle is 12 degrees. The tension  $T$  (in N) in the back muscle is:

- A) 460      B) 2117      C) 0  
 D) 722      E) 2164




28) The horizontal component of the force  $R$  ( $\equiv R_x$ ) exerted by the pivot (sacrum) along the spine (in N)?

- A) 2117      B) 450      C) 0  
 D) 2164      E) 1667

Dr. Ahmad S. Y.

## Department Of Physics

1) The density of water is  $1.0 \text{ g/cm}^3$ . If  $h = 20 \text{ cm}$ , the density of the oil in the left column of the U-tube shown below is:



- A)  $0.20 \text{ g/cm}^3$
- B)  $0.90 \text{ g/cm}^3$**
- C)  $1.0 \text{ g/cm}^3$
- D)  $1.3 \text{ g/cm}^3$
- E)  $5.0 \text{ g/cm}^3$

2) One piston in a hydraulic lift has an area that is twice the area of the other. When the pressure at the smaller piston is increased by  $\Delta p$  the pressure at the larger piston:

- A) increases by  $2\Delta p$
- B) increases by  $\Delta p/2$
- C) increases by  $\Delta p$**
- D) increases by  $4\Delta p$
- E) does not change

3) A boat floating in fresh water displaces 16,000 N of water. How many newtons of salt water would it displace if it floats in salt water of specific gravity 1.10?

- A) 12,800 N
- B) 14,400 N
- C) 16,000 N**
- D) 17,600 N
- E) 19,200 N

4) An object hangs from a spring balance. The balance indicates 30 N in air, 20 N when the object is submerged in water. What does the balance indicate when the object is submerged in liquid with a density that is half of water?

- A) 20 N
- B) 25 N**
- C) 30 N
- D) 35 N
- E) 40 N

5) The dimensions of a wooden raft (density =  $150 \text{ kg/m}^3$ ) are  $3.0 \text{ m} \times 3.0 \text{ m} \times 1.0 \text{ m}$ . What maximum load can it carry in sea water (density =  $1020 \text{ kg/m}^3$ )?

- A) 1350 kg
- B) 7830 kg
- C) 9200 kg
- D) 19,500 kg
- E) 24,300 kg

6) A lawn sprinkler is made of a 1.0 cm diameter garden hose with one end closed and 25 holes, each with a diameter of 0.050 cm, cut near the closed end. If water flows at 2.0 m/s in the hose, the speed of the water leaving a hole is:

- A) 2.0 m/s
- B) 32 m/s
- C) 40 m/s
- D) 600 m/s
- E) 800 m/s

7) Water is streaming downward from a faucet opening with an area of  $3.0 \times 10^{-5} \text{ m}^2$ . It leaves the faucet with a speed of 5.0 m/s. The cross sectional area of the stream 0.50 m below the faucet is:

- A)  $1.5 \times 10^{-5} \text{ m}^2$
- B)  $2.0 \times 10^{-5} \text{ m}^2$
- C)  $2.5 \times 10^{-5} \text{ m}^2$
- D)  $3.0 \times 10^{-5} \text{ m}^2$
- E)  $3.5 \times 10^{-5} \text{ m}^2$

8) A fluid of density  $9.1 \times 10^2 \text{ kg/m}^3$  is flowing through a tube at a speed of 5.3 m/s. What is the kinetic energy density of the fluid?

- A) cannot be calculated without knowing the pressure
- B) cannot be calculated without knowing the elevation
- C)  $4.8 \times 10^3 \text{ J/m}^3$
- D)  $1.3 \times 10^4 \text{ J/m}^3$
- E)  $2.5 \times 10^6 \text{ J/m}^3$

9) Water (density =  $1.0 \times 10^3 \text{ kg/m}^3$ ) flows downhill through a pipe of diameter 1.5 cm. Its speed at the top of the hill is 7.2 m/s. If the hill is 9.5 m high, what is the gravitational potential energy density of the water at the top of the hill relative to the bottom?

- A) cannot be calculated without knowing the pressure
- B)  $120 \text{ J/m}^3$
- C)  $7.2 \times 10^3 \text{ J/m}^3$
- D)  $9.5 \times 10^3 \text{ J/m}^3$
- E)  $9.3 \times 10^4 \text{ J/m}^3$

10) Water (density =  $1.0 \times 10^3 \text{ kg/m}^3$ ) flows through a horizontal tapered pipe. At the wide end its speed is 4.0 m/s. The difference in pressure between the two ends is  $4.5 \times 10^3 \text{ Pa}$ . The speed of the water at the narrow end is:

- A) 2.6 m/s
- B) 3.2 m/s
- C) 4.0 m/s
- D) 4.5 m/s
- E) 5.0 m/s

11) A large tank filled with water has two holes in the bottom, one with twice the radius of the other. In steady flow the speed of water leaving the larger hole is \_\_\_\_\_ the speed of the water leaving the smaller.

- A) twice
- B) four times
- C) half
- D) one-fourth
- E) the same as

12) Some species of whales can dive to depths of one kilometer. What is the total pressure they experience at this depth? ( $\rho_{\text{sea}} = 1020 \text{ kg/m}^3$  and  $1.01 \times 10^5 \text{ N/m}^2 = 1 \text{ ATM}$ .)

- a. 9.00 ATM
- b. 90.0 ATM
- c. 100 ATM
- d. 111 ATM
- e. 130 ATM

13) Water is flowing at 4.0 m/s in a circular pipe. If the diameter of the pipe decreases to 1/2 its former value, what is the velocity of the water downstream?

- a. 1.0 m/s
- b. 2.0 m/s
- c. 8.0 m/s
- d. 16 m/s
- e. 4.0 m/s

14) What is the net force inward acting on a spherical bathysphere of diameter 2.00 m at an ocean depth of 1000 m? (The pressure inside the bathysphere is, hopefully, 1 ATM.)

$\rho_{\text{sea water}} = 1.02 \times 10^3 \text{ kg/m}^3$ .

- a.  $1.26 \times 10^4 \text{ N}$
- b.  $1.26 \times 10^6 \text{ N}$
- c.  $1.26 \times 10^8 \text{ N}$
- d.  $1.26 \times 10^{10} \text{ N}$
- e.  $1.26 \times 10^2 \text{ N}$

15) How much power is theoretically available from a mass flow of 1 000 kg/s of water when it falls a vertical distance of 100 meters?

- a. 980 kW
- b. 98 kW
- c. 4 900 W
- d. 980 W

Dr. Ahmad Masadeh

e. 9 600 W


16) A cubical box, 5.00 cm on each side, is immersed in a fluid. The gauge pressure at the top surface of the box is 594 Pa and the gauge pressure on the bottom surface is 1133 Pa. What is the density of the fluid?

- A) 1000 kg/m<sup>3</sup>
- B) 1100 kg/m<sup>3</sup>
- C) 1220 kg/m<sup>3</sup>
- D) 2340 kg/m<sup>3</sup>
- E) 12,000 kg/m<sup>3</sup>

17) The weight of a car of mass  $1.20 \times 10^3$  kg is supported equally by the four tires, which are inflated to the same gauge pressure. What gauge pressure in the tires is required so the area of contact of each tire with the road is  $1.00 \times 10^2$  cm<sup>2</sup>? (1 atm =  $1.01 \times 10^5$  Pa.)


- A)  $11.6 \times 10^5$  Pa
- B)  $11.6 \times 10^4$  Pa
- C)  $2.94 \times 10^5$  Pa
- D)  $2.94 \times 10^4$  Pa
- E)  $2.94 \times 10^3$  Pa

18) In the figure, an open tank contains a layer of oil floating on top of a layer of water (of density 1000 kg/m<sup>3</sup>) that is 3.0 m thick, as shown. What must be the thickness of the oil layer if the gauge pressure at the bottom of the tank is to be  $5.0 \times 10^4$  Pa? The density of the oil is 510 kg/m<sup>3</sup>.



Answer: 4.12 m

19) The two water reservoirs shown in the figure are open to the atmosphere, and the water has density  $1000 \text{ kg/m}^3$ . The manometer contains incompressible mercury with a density of  $13,600 \text{ kg/m}^3$ . What is the difference in elevation  $h$  if the manometer reading  $m$  is  $25.0 \text{ cm}$ ?



- A) 1.58 m
- B) 4.20 m
- C) 3.75 m
- D) 3.40 m
- E) 3.15 m

20) A board that is  $20.0 \text{ cm}$  wide,  $5.00 \text{ cm}$  thick, and  $3.00 \text{ m}$  long has a density  $350 \text{ kg/m}^3$ . The board is floating partially submerged in water of density  $1000 \text{ kg/m}^3$ . What fraction of the volume of the board is above the surface of the water?

- A) 0.350
- B) 0.650
- C) zero
- D) 0.200
- E) The answer depends on which edge of the board is vertical.

21) A person who weighs  $550 \text{ N}$  empties her lungs as much as possible and is then completely immersed in water (of density  $1000 \text{ kg/m}^3$ ) while suspended from a harness. Her apparent weight is now  $21.2 \text{ N}$ . What is her density?

- A)  $1050 \text{ kg/m}^3$
- B)  $1040 \text{ kg/m}^3$
- C)  $1030 \text{ kg/m}^3$
- D)  $960 \text{ kg/m}^3$
- E)  $56.1 \text{ kg/m}^3$

22) A  $7.8\text{-kg}$  solid sphere, made of metal whose density is  $2500 \text{ kg/m}^3$ , is suspended by a cord. When the sphere is immersed in water (of density  $1000 \text{ kg/m}^3$ ), what is the


tension in the cord?

- A) 46 N
- B) 61 N
- C) 76 N
- D) 92 N
- E) 110 N

23) Water flowing through a pipe suddenly comes to a section of pipe where the pipe diameter decreases to 86% of its previous value. If the speed of the water in the larger section of the pipe was 36m/s, what is its speed in this smaller section?

- A) 49 m/s
- B) 42 m/s
- C) 31 m/s
- D) 27 m/s

24) Water flows in the horizontal pipe shown in the figure. At point A the area is 25.0 cm<sup>2</sup> and the speed of the water is 2.00 m/s. At B the area is 16.0 cm<sup>2</sup>. The fluid in the manometer is mercury, which has a density of 13,600 kg/m<sup>3</sup>. We can treat water as an ideal fluid having a density of 1000 kg/m<sup>3</sup>. What is the manometer reading  $h$ ?



- A) 0.546 cm
- B) 1.31 cm
- C) 2.81 cm
- D) 2.16 cm
- E) 3.36 cm



**The University Of Jordan**

**Faculty of Science**

**Department Of Physics**

1) A 100-kg box rolls down a  $20^\circ$  incline. A man tries to keep it from accelerating, and manages to keep its acceleration to  $1.2 \text{ m/s}^2$ . If the box rolls 5 m, what is the net work done on it by all the forces acting on it?

- A) 60 J
- B) 100 J
- C) 600 J
- D) 1000 J
- E) 4900 J

2) Two objects with masses,  $m_1$  and  $m_2$ , have the same kinetic energy and are both moving to the right. The same constant force  $\vec{F}$  is applied to the left to both masses. If  $m_1 = 4m_2$ , the ratio of the stopping distance of  $m_1$  to that of  $m_2$  is: A)

- 1:4
- B) 4:1
- C) 1:2
- D) 2:1
- E) 1:1

3) A 4-kg cart starts up an incline with a speed of 3 m/s and comes to rest 2 m up the incline. The total work done on the cart is:

- A) -6 J
- B) -8 J
- C) -12 J
- D) -18 J
- E) impossible to calculate without knowing the coefficient of kinetic friction


4) A 50-N force is the only force acting on a 2-kg crate that starts from rest. When the force has been acting for 2 s the rate at which it is doing work is:

- A) 100 W
- B) 1000 W
- C) 2500 W
- D) 5000 W
- E) 63000 W

5) A 6.0-kg block is released from rest 80 m above the ground. When it has fallen 60 m its kinetic energy is approximately:


- A) 4700 J
- B) 3500 J
- C) 1200 J
- D) 120 J
- E) 60 J

6) For a block of mass  $m$  to slide without friction up the rise of height  $h$  shown, it must have a minimum initial kinetic energy of:



- A)  $gh$
- B)  $mgh$
- C)  $gh/2$
- D)  $mgh/2$
- E)  $2mgh$

7) A simple pendulum consists of a 2.0 kg mass attached to a string. It is released from rest at X as shown. Its speed at the lowest point Y is:




- A) 1.9 m/s
- B) 3.7 m/s
- C) 4.4 m/s
- D) 6.0 m/s
- E) 36 m/s

8) A 2.2-kg block starts from rest on a rough inclined plane that makes an angle of  $25^\circ$  with the horizontal. The coefficient of kinetic friction is 0.25. As the block goes 2.0 m down the plane, the mechanical energy of the whole system changes by:

- A) 0 J
- B) -9.8 J
- C) 9.8 J
- D) -18 J
- E) 18 J

12) A 4.0-kg block is lowered down a  $37^\circ$  incline a distance of 5.0 m from point A to point B. A horizontal force ( $F = 10$  N) is applied to the block between A and B as shown in the figure. The kinetic energy of the block at A is 10 J and at B it is 20 J. How much work is done on the block by the force of friction between A and B?



- a. -58 J
- b. -53 J
- c. -68 J
- d. -63 J
- e. -47 J


13) A 0.60-kg object is suspended from the ceiling at the end of a 2.0-m string. When pulled to the side and released, it has a speed of 4.0 m/s at the lowest point of its path. What maximum angle does the string make with the vertical as the object swings up?

- a.  $61^\circ$
- b.  $54^\circ$
- c.  $69^\circ$
- d.  $77^\circ$
- e.  $47^\circ$

14) A 2.0-kg mass swings at the end of a light string (length = 3.0 m). Its speed at the lowest point on its circular path is 6.0 m/s. What is its kinetic energy at an instant when the string makes an angle of  $50^\circ$  with the vertical?


- a. 21 J
- b. 15 J
- c. 28 J
- d. 36 J
- e. 23 J

9) A rod is pivoted about its center. A 5-N force is applied 4 m from the pivot and another 5-N force is applied 2 m from the pivot, as shown. The magnitude of the total torque about the pivot is:




- A) 0 N·m
- B) 5.0 N·m
- C) 8.7 N·m
- D) 15 N·m
- E) 26 N·m

10) A uniform rod AB is 1.2 m long and weighs 16 N. It is suspended by strings AC and BD as shown. A block P weighing 96 N is attached at E, 0.30 m from A. The magnitude of the tension force in the string BD is:



- A) 8.0 N
- B) 24 N
- C) 32 N
- D) 48 N
- E) 80 N

11) An 80-N uniform rod leans against a frictionless wall as shown. The torque (about point P) applied to the rod by the wall is:



- A) 40 N·m
- B) 60 N·m
- C) 120 N·m
- D) 160 N·m
- E) 240 N·m

15) The same force  $F$  is applied horizontally to bodies 1, 2, 3 and 4, of masses  $m$ ,  $2m$ ,  $3m$  and  $4m$ , initially at rest and on a frictionless surface, until each body has traveled distance  $d$ . The correct listing of the magnitudes of the velocities of the bodies,  $v_1$ ,  $v_2$ ,  $v_3$ , and  $v_4$  is

- a.  $v_4 = \sqrt{\frac{4}{3}} v_3 = \sqrt{\frac{3}{2}} v_2 = 2v_1$ .
- b.  $v_4 = v_2 > v_3 = v_1$ .
- c.  $v_1 = \sqrt{2} v_2 = \sqrt{3} v_3 = 2v_4$ .
- d.  $v_1 = 2v_2 = 3v_3 = 4v_4$ .
- e.  $v_4 = \frac{3}{4} v_3 = \frac{2}{3} v_2 = \frac{1}{2} v_1$ .

16) A 3.0-kg block is on a frictionless horizontal surface. The block is at rest when, at  $t = 0$ , a force (magnitude  $P = 2.0$  N) acting at an angle of  $22^\circ$  above the horizontal is applied to the block. At what rate is the force  $P$  doing work at  $t = 2.0$  s?

- a. 2.3 W
- b. 2.0 W
- c. 1.4 W
- d. 1.7 W
- e. 1.2 W

17) A 3.0-kg block is on a horizontal surface. The block is at rest when, at  $t = 0$ , a force (magnitude  $P = 12$  N) acting parallel to the surface is applied to the block causing it to accelerate. The coefficient of kinetic friction between the block and the surface is 0.20. At what rate is the force  $P$  doing work on the block at  $t = 2.0$  s?

- a. 54 W
- b. 49 W
- c. 44 W
- d. 59 W
- e. 24 W


18) A crane lifts a 425 kg steel beam vertically a distance of 117 m. How much work does the crane do on the beam if the beam accelerates upward at  $1.8 \text{ m/s}^2$ ? Neglect frictional forces.

- A)  $5.8 \times 10^5$  J
- B)  $3.4 \times 10^5$  J
- C)  $4.0 \times 10^5$  J
- D)  $4.9 \times 10^5$  J

19) A 1000.0 kg car is moving at 15 km/h. If a 2000.0 kg truck has 18 times the kinetic energy of the car, how fast is the truck moving?

- A) 45 km/h
- B) 63 km/h
- C) 54 km/h
- D) 36 km/h

20) In the figure, two boxes, each of mass 24 kg, are at rest and connected as shown. The coefficient of kinetic friction between the inclined surface and the box is 0.31. Find the speed of the boxes just after they have moved 1.6 m. **Answer: 1.91 m/s**




21) A car needs to generate 75.0 hp in order to maintain a constant velocity of 27.3 m/s on a flat road. What is the magnitude of the total resistive force acting on the car (due to friction, air resistance, etc.)? (1 hp = 746 W)

- A)  $2.05 \times 10^3$  N
- B) 2.75 N
- C)  $1.03 \times 10^3$  N
- D)  $2.87 \times 10^3$  N

22) How long will it take a 7.08 hp motor to lift a 250 kg beam directly upward at constant velocity from the ground to a height of 45.0 m? Assume frictional forces are negligible. (1 hp = 746 W)

- A) 20.9 s
- B)  $1.56 \times 10^4$  s
- C)  $2.18 \times 10^4$  s
- D) 39.7 s

23) A roller coaster of mass 80.0 kg is moving with a speed of 20.0 m/s at position A as shown in the figure. The vertical height above ground level at position A is 200 m. Neglect friction.



(a) What is the total mechanical energy of the roller coaster at point B?

Answer:  $1.73 \times 10^5 \text{ J}$

(b) What is the speed of the roller coaster at point C?


Answer: 34.4 m/s

24) In the figure, a block of mass  $m$  is moving along the horizontal frictionless surface with a speed of 5.70 m/s. If the slope is  $11.0^\circ$  and the coefficient of kinetic friction between the block and the incline is 0.260, how far does the block travel up the incline?




Answer: 3.72 m

25) A 10.0-kg uniform ladder that is 2.50 m long is placed against a smooth vertical wall and reaches to a height of 2.10 m, as shown in the figure. The base of the ladder rests on a rough horizontal floor whose coefficient of static friction with the ladder is 0.800. An 80.0-kg bucket of concrete is suspended from the top rung of the ladder, right next to the wall, as shown in the figure. What is the magnitude of the friction force that the floor exerts on the ladder?



- A) 538 N
- B) 706 N
- C) 1290 N
- D) 833 N
- E) 601 N


26) A solid uniform brick is placed on a sheet of wood. When one end of the sheet is raised (see figure), you observe that the maximum that the angle  $\theta$  can be without tipping over the brick is  $49.6^\circ$ . There is enough friction to prevent the brick from sliding. What is the width  $w$  of the brick?



- A) 5.18 cm
- B) 6.09 cm
- C) 6.81 cm
- D) 9.40 cm
- E) 10.5 cm



Person X pushes twice as hard against a stationary brick wall as person Y. Which one of the following statements is correct?

- A. Both do positive work, but person X does four times the work of person Y.
- B. Both do positive work, but person X does twice the work of person Y.
- C. Both do the same amount of positive work.
- D. Each one of them does zero work 
- E. Both do positive work, but person X does one-half the work of person Y.

What is the average power output (in  $W$ ) of a  $60.0\text{-kg}$  athlete when, in  $8.00\text{ s}$ , he runs up a flight of stairs that is  $10.0\text{-m}$  high at constant speed?

- A.  $75.0$
- B.  $735$
- C.  $4800$
- D.  $48$
- E.  $600$




A 60-kg skier starts from rest from the top of a 50-m high slope. If the work done by friction is  $-6.0$  kJ, what is the speed (in m/s) of the skier on reaching the bottom of the slope?

- A. 17
- B. 24
- C. 28
- D. 31
- E. 57



A bead is moving with a speed of  $20 \text{ m/s}$  at position A on the track shown in the figure. This track is friction-free. What is the speed (in  $\text{m/s}$ ) of the bead at point C?



- A. 0
- B. 34
- C. 69
- D. 20
- E. We cannot solve this problem without knowing the mass of the bead.

A 4.0 kg object is moving with speed 2.0 m/s. A 1.0 kg object is moving with speed 4.0 m/s. Both objects encounter the same constant braking force, and are brought to rest. Which object travels the greater distance before stopping?

- A. the 4.0 kg object
- B. the 1.0 kg object
- C. both objects travel the same distance ✓
- D. answer cannot be determined from the information given
- E. The 4 kg object travels twice the distance covered by the 1 kg object

A truck has four times the mass of a car and is moving with twice the speed of the car. If  $K_t$  and  $K_c$  refer to the kinetic energies of truck and car respectively, it is correct to say that

- A.  $K_t = 16K_c$
- B.  $K_t = 4K_c$
- C.  $K_t = 2K_c$
- D.  $K_t = K_c$
- E.  $K_t = K_c$



A 35-N bucket of water is lifted vertically 3.0 m and then returned to its original position. How much work (in J) did gravity do on the bucket during this process?

- A. 180
- B. 90
- C. 0
- D. 900
- E. 45



When a car of mass 1167 kg accelerates from 10.0 m/s to some final speed,  $4.00 \times 10^5$  J of work are done. Find this final speed (in m/s).

A. 28.0

B. 22.4

C. 25.2


D. 30.8

E. 16.7






A person carries a 2.00-N object through the path shown in the figure, starting at point A and ending at point B. The total time from A to B is 6.75 min. How much work did gravity do on the object between A and B?



- a. 30.0
- b. -30
- c. -56
- d. -36
- e. 0



A stone can slide down one of four different frictionless ramps, as shown in the figure. For which ramp will the speed of the ball be the greatest at the bottom?



- A. Ramp X
- B. Ramp Y
- C. Ramp Z
- D. The speed of the ball will be the same for all ramps ✓
- E. More information is needed to answer the question

Which of the following statements is CORRECT?

Select one:

- a. An object can accelerate even when the resultant force acting on it is zero.
- b. When you walk forward without skidding, the static friction is the force that caused you to move. ✓
- c. Weight is a scalar quantity.
- d. The normal force is the reaction force to the weight of an object.
- e. Acceleration is always in opposite direction to the resultant force.

The kinetic friction force that a horizontal surface exerts on a 60.0-kg object is 50.0 N. If the initial speed of the object is 25.0 m/s, what distance (in m) will it slide before coming to a stop?

- A. 15.0
- B. 30.0
- C. 375
- D. 750
- E. 855



A stone initially moving at  $8.0 \text{ m/s}$  on a level surface comes to rest due to friction after it travels  $11 \text{ m}$ . What is the coefficient of kinetic friction between the stone and the surface?

- A. 0.13
- B. 0.50
- C. 0.30
- D. 0.43
- E. 0.80



A student starts from the origin at  $t=0$  s. He moved along the positive  $x$ -direction for 6.0 m. Then he moved along the negative  $x$ -direction a distance of 3.0 m. If the total time of his motion is 6.0 s, then his average speed (in m/s) is:

- a. 2.0
- b. 0
- c. 3.0
- d. 1.0
- e. 1.5



The correct answer is: 1.5

The velocities (in m/s) of cars A and B are given at equal time intervals.


|        |    |    |    |    |
|--------|----|----|----|----|
| Car A: | 20 | 20 | 20 | 20 |
| Car B: | 1  | 3  | 5  | 7  |

Which of the following statements is correct?

- a. Neither car accelerates
- b. Car A has variable velocity
- c. Car A does not accelerate and car B accelerates. ✓
- d. Car B is moving along the negative x-direction
- e. Car A has larger acceleration than car B

The correct answer is: Car A does not accelerate and car B accelerates.

Vectors  $A = 3.0$  m and  $B = 8.0$  m are represented as shown in the figure. What is the magnitude (in m) of the resultant ?




- a. 7.0
- b. 3.9
- c. 11.0
- d. 4.2
- e. 8.4



The correct answer is: 7.0



Which of the following can be used as a conversion factor to write m/s as mi/h? (1 mi = 1609 m)

- a.  $(1609/3600)$  mi/h
- b.  $(3600/1609)$  mi/h 
- c.  $(1609/3600)$  h/mi
- d.  $(3600/1609)$  h/mi
- e. 3600 s/h

The correct answer is:  
 $(3600/1609)$  mi/h

Which of the following statements is correct?

- a. If an object moves, its average velocity can NEVER be zero.
- b. A car moving at constant velocity has non zero acceleration.
- c. Average velocity depends on distance
- d. If an object moves its average velocity can be zero, but its average speed must be greater than zero. ✓
- e. Average speed depends on displacement

The position of a runner is  $x = 2.0$  m at  $t = 1.0$  s . At  $t = 3.0$  s the new position of the runner is  $x = 5.0$  m. The average velocity (in m/s) of the runner over the time interval from 1.0 to 3.0 s is:

- a. 1.5
- b. 3.0
- c. 0
- d. 1.0
- e. 6.0



The density of gold is 19000 kg/m<sup>3</sup>. The density of gold in gram/cm<sup>3</sup> is:

- a. 1
- b. 19
- c. 0.19
- d. 1900
- e. 190



The correct answer is: 19

A car moves from point A to point B at a speed of 25 km/h. It then moved from point B back to point A at a speed of 20 km/h. The average speed (in km/h) of the car is:

- a. 22.5
- b. 22.2
- c. 23.7
- d. 11.1
- e. 21.9



The correct answer is: 22.2

The position of a car is given by the equation  $x = A + B t^2$ . The dimensions of the constants A and B, respectively, are:

- a.  $T^2/L$
- b.  $L^2$  and  $L T^2$
- c.  $L^2$  and  $L/T^4$
- d.  $L$  and  $L/T^2$
- e.  $T^2$



The correct answer is:  $L$  and  $L/T^2$

A car is moving at 35 km/h.  
The speed of the car in m/s is:

- a. 35
- b. 9.7
- c. 126
- d. 75
- e. 22




The correct answer is: 9.7

If a car is moving to the left with constant velocity, one can conclude that

- A. there must be no forces applied to the car.
- B. there is exactly one force applied to the car.
- C. The net force applied to the car must be to the right
- D. the net force applied to the car is directed to the left.
- E. the net force applied to the car is zero. ✓



Three blocks, light connecting ropes, and a light frictionless pulley comprise a system, as shown in the figure. An external force of magnitude  $P$  is applied downward on block A, causing block A to accelerate downward at a constant  $2.5 \text{ m/s}^2$ . The tension in the rope connecting block B and block C is equal to  $60 \text{ N}$ . The mass (in kg) of block C is:



- A. 18.0
- B. 3.5
- C. 9.8
- D. 4.9
- E. 6.0




A trolley is carrying a 20.0-kg box along a level road. The coefficient of static friction between the box and the floor of the trolley is 0.400, and the coefficient of kinetic friction is 0.300. What is the maximum acceleration (in  $\text{m/s}^2$ ) that the trolley can have if the box is to move with the trolley without sliding?

- A. 7.40
- B. 196
- C. 3.92
- D. 8.00
- E. 78.5



In the figure the 9.0-kg block is on a smooth horizontal table. The surfaces of the 12-kg block are rough, with  $\mu_k = 0.30$  between the 12-kg and 9.0-kg blocks. The mass  $M=5.0$  and accelerates downwards. The acceleration of mass  $M$  (in  $\text{m/s}^2$ ).




- A. 5.7
- B. 6.2
- C. 1.9
- D. 3.1
- E. 0.98



An object can remain at rest

- A. ONLY when there are ✗  
no forces at all acting  
on it.
- B. when the net force acting  
on it is zero
- C. when the net force acting  
on it is a nonzero  
constant.
- D. when there is only one  
force acting on it.
- E. Only when no frictional  
forces acting on it

A box of mass of  $m_1 = 10$  kg collides with a box of mass  $m_2 = 2$  kg. Which of the following statements is correct?

- A.  $m_1$  acts with a force on  $m_2$  but  $m_2$  does not act with a force on  $m_1$  because it is small
- B. No force is exchanged between  $m_1$  and  $m_2$
- C. The force of  $m_1$  on  $m_2$  is five times larger than the force of  $m_2$  on  $m_1$
- D. The force of  $m_2$  on  $m_1$   is equal to the force of  $m_1$  on  $m_2$
- E. The force of  $m_2$  on  $m_1$  is larger than the force of  $m_1$  on  $m_2$

A person is using a rope to lower a 5.0-N bucket into a well with a constant speed of 2.0 m/s. What is the magnitude of the force exerted by the rope on the bucket?

- A. 10
- B. 5
- C. 0
- D. 2
- E. 49



A 60-kg person is in an elevator that is moving down and accelerating at  $2 \text{ m/s}^2$ . His apparent weight (in N) is: (take  $g=9.8 \text{ m/s}^2$ )

- A. 468
- B. 590
- C. 588
- D. zero
- E. 660




A 1200-kg car is pulling a 500-kg trailer along level ground. Friction of the road on the trailer is negligible. The car accelerates with an acceleration of  $1.3 \text{ m/s}^2$ . What is the force exerted by the car on the trailer?

- A. 550
- B. 600
- C. 700
- D. 650
- E. 300






A man raises a 1000-N piano at a constant speed using a very light rope in a frictionless pulley system, as shown in the figure. With what force is the man pulling down on the rope?



- A. 250
- B. 1000
- C. 500
- D. 2000
- E. 1500




The figure shows a block of mass  $M$  hanging at rest. The light wire fastened to the wall is horizontal and has a tension of 38 N. The wire fastened to the ceiling is also very light, has a tension of 59 N and makes an angle  $\theta$  with the ceiling. Find the angle  $\theta$  (in degrees).




- A. 40
- B. 65
- C. 45
- D. 33
- E. 50



In the figure the surface of the table is smooth and the system is released from rest. What is the magnitude of the acceleration of the 10-kg block (in  $\text{m/s}^2$ ) when the system is released from rest?



Two masses  $m_1 = 2.0 \text{ kg}$  and  $m_2 = 4.0 \text{ kg}$  are connected by a light inextensible string as shown in the figure. The system is pulled along a frictionless surface by a force  $F = 18 \text{ N}$ . The value of the tension  $T$  (in N) is:



Select one:

- a. 24.0
- b. 3.0
- c. 6.0 ✓
- d. 12.0
- e. 18.0

In the figure mass  $M = 4.0$  kg and mass  $m = 2.0$  kg. The ground surface is frictionless, while the coefficient of static friction between the two masses is 0.30. Find the maximum value of  $F$  (in N) such that mass  $m$  moves with mass  $M$  without sliding.



A diagram showing two rectangular blocks stacked on a horizontal surface. The top block is orange and labeled with the lowercase letter 'm'. The bottom block is blue and labeled with the uppercase letter 'M'. A thick black horizontal line is positioned below the blue block, representing the surface it rests on. A thin black horizontal line is positioned to the right of the blue block, extending from its right edge.

$m$

$M$



Select one:

a. 25.9

b. 3.2

c. 17.6 ✓

Well Done!

d. 11.8

e. 5.9

What force (in N) is needed to stop a 1000-kg car moving at 25 m/s during a time interval of 10 seconds?

Select one:

- a. 400
- b. 500
- c. 250
- d. 2000
- e. 2500 ✓

A 2.0-kg block is on the verge of sliding down a rough inclined plane that makes an angle of 40 degrees with the horizontal. The coefficient of static friction  $\mu_s$  is:

Select one:

A. 0.50

B. 0

C. 0.84 ✓ Well done!

D. 0.64

E. 0.77

A block of mass  $m = 4.0 \text{ kg}$  slides down a  $35^\circ$  incline when a force of  $F = 10 \text{ N}$  is applied upward parallel to the incline. If the coefficient of kinetic friction between the block and the incline is  $0.2$ , find the acceleration (in  $\text{m/s}^2$ ) of the block as it moves down the inclined plane:

Select one:


- a. 3.1
- b. 4.0
- c. 0.44
- d. 2.7
- e. 1.5 ✓

A force accelerates a body of mass  $M$ . The same force applied to a second body produces three times the acceleration. The mass of the second body will be:

Select one:

- a.  $2M$
- b.  $M/3$  ✓ Well Done!
- c.  $M/2$
- d.  $9M$
- e.  $3M$

Two objects each of mass  $M$  are connected by a light inextensible cord. The system is attached by another cord to the ceiling of an elevator that is accelerating upward at  $2 \text{ m/s}^2$ , the ratio of the tensions  $T_1/T_2$  is:





Select one:

a. 2

b. 1


c.  $\frac{5}{3}$

d.  $\frac{3}{2}$

e.  $\frac{1}{2}$  ✓

Well Done!

In the figure, mass  $M = 25 \text{ kg}$  is in static equilibrium. The value of the tension  $T_1$  (in Newton) is:



Select one:


- a. 245
- b. 147.4
- c. 195.7 ✓
- d. 201.2
- e. 0

A 50-N crate sits on a horizontal floor where the coefficient of static friction between the crate and the floor is 0.50. A 20-N force is applied to the crate acting to the right. What is the resulting static friction force (in N) acting on the scale?

Select one:

- a. 20 to the left. ✓
- b. 25 to the left.
- c. 20 to the right.
- d. 0
- e. 25 to the right.

The figure below shows a container filled with water to the height shown. When we compare the pressure at a ( $P_a$ ) to the pressure at b ( $P_b$ ), we find that



Select one:

- a.  $P_a = 2 P_b$
- b.  $P_a = P_b/2$
- c.  $P_a = P_b$  ✓
- d.  $P_a = 4 P_b$

The pressure inside a plane is maintained at 1 atm ( $1.013 \times 10^5 \text{ N/m}^2$ ). What is the net force (in N) exerted on a 10 m x 2.0 m cabin door if the outside pressure (at 10 km height) is 0.3 atm?


Select one:

- a.  $1.42 \times 10^5$  ✓
- b. 0
- c.  $1.013 \times 10^5$
- d.  $0.6 \times 10^5$
- e.  $2.03 \times 10^5$

The correct answer is:  $1.42 \times 10^5$

Two identical blocks of ice float in water as shown. Which of the following statements is

correct?



Select one:

- a. block A displaces a greater volume of water since the pressure acts on a smaller bottom area
- b. block B displaces a greater volume of water since the pressure is less on its bottom
- c. the two blocks displace equal volumes of water since they have the same weight ✓
- d. block A displaces a greater volume of water since its submerged end is lower in the water
- e. block B displaces a greater volume of water since its submerged end has a greater area




A table-tennis ball has a diameter of 3.80 cm and a density of  $84 \text{ kg/m}^3$ . What force is required to hold it completely submerged under water? (water density =  $1000 \text{ kg/m}^3$ )

Select one:

- a. 0
- b. 0.281
- c. 0.024
- d. 0.258 ✓
- e. 0.018

The correct answer is: 0.258

The vessels shown below all contain water to the same height. Rank them according to the pressure exerted by the water on the vessel bottoms, least to greatest.



Select one:


- a. 2, 4, 3, 4
- b. All the pressures are the same. ✓
- c. 4, 3, 2, 1
- d. 1, 2, 3, 4
- e. 1, 2, 4, 3

A hydraulic press has one piston of diameter 2.0 cm and the other piston of diameter 8.0 cm. What force (in N) must be applied to the smaller piston to obtain a force of 1600 N at the larger piston.

Select one:

- a. 100 ✓
- b. 26000
- c. 1600
- d. 400
- e. 6400

A cubical block of wood of side length 10.0 cm floats in equilibrium at the interface between oil and water with its lower surface 2.00 cm below the water surface as shown. The density of the oil is  $750 \text{ kg/m}^3$ . Calculate the mass (in Kg) of the block. (density of water is  $1000 \text{ kg/m}^3$ )



Select one:

- a. 0.80 ✓
- b. 0.60
- c. 2.00
- d. 1.40
- e. 0.20

A small boat is 4.0 m wide and 6.0 m long. When a truck is loaded onto the boat, the boat sinks and additional 5.0 cm into the river. What is the weight (in N) of the truck? (assume density of sea water to be  $1025 \text{ kg/m}^3$ ).

Select one:

- a. 12054 ✓
- b. 241080
- c. 324235
- d. 15432
- e. 23456


A plastic sphere floats in water with 50% of its volume below the water surface. What is the density (in  $\text{Kg/m}^3$ ) of the plastic sphere? (water density =  $1000 \text{ kg/m}^3$ )

Select one:

- a. 100
- b. 250
- c. 1000
- d. 50
- e. 500 ✓

In the figure, the larger area has a diameter  $D_1 = 2.50$  cm. Water flows to the right at a rate of  $180 \times 10^{-4}$  m<sup>3</sup>/s. Determine the diameter  $D_2$  (in cm) of the smaller area. (density of water is

1000 kg/m<sup>3</sup>)



Select one:

- a. 2.6
- b. 1.47 ✓
- c. 2.12
- d. 1.78
- e. 1.89

An object is moving along the positive x-direction with an acceleration of  $-3 \text{ m/s}^2$ . Which of the following statements is correct?

Select one:

- a. The object will always be moving in the the positive x-direction.
- b. The object will accelerate
- c. The speed of the object will decrease. ✓
- d. The object will never reverse its direction of motion.
- e. The speed of the object will increase

The correct answer is: The speed of the object will decrease.

Correct





Determine the stopping distance (in m) for an automobile moving with an initial speed of 25 m/s, if it decelerates at  $2.5 \text{ m/s}^2$  and the driver's reaction time is 0.4 s.

Select one:

- a. 10
- b. 125
- c. 135 ✓ Well done!
- d. 100
- e. 625

The correct answer is: 135

Correct

Marks for this submission: 2.0/2.0.

A stone is projected vertically upwards. Which of the following statements is WRONG?

Select one:

Correct  
Marks for this submission: 2.0/2.0.

A stone is projected vertically upwards. Which of the following statements is WRONG?

Select one:

- a. At maximum height its acceleration is zero ✓ Well done
- b. As it moves up its speed decreases
- c. Its acceleration is always  $9.8 \text{ m/s}^2$  towards the center of the earth.
- d. As it moves down its speed increases
- e. When it reverses its direction of motion it has zero velocity

The correct answer is: At maximum height its acceleration is zero

Correct

Marks for this submission: 2.0/2.0.

A rocket rises vertically from rest with an acceleration of  $3.0 \text{ m/s}^2$  until it runs out of fuel at a height of 600 m. After this it is in free fall motion. How long (in s) (from the moment the fuel runs out) will it take the rocket to reach the ground?

A rocket rises vertically from rest with an acceleration of  $3.0 \text{ m/s}^2$  until it runs out of fuel at a height of 600 m. After this it is in free fall motion. How long (in s) (from the moment the fuel runs out) will it take the rocket to reach the ground?

Select one:

- a. 18.8 ✓ Well done!
- b. 23.5
- c. 33.1
- d. 6.5
- e. 60.0

The correct answer is: 18.8

**Correct**

Marks for this submission: 2.0/2.0.

An object is thrown vertically upward from the top of a 30 m high building with an initial speed of 20 m/s. The average velocity (in m/s) during the time interval  $t=0$  to  $t=5$  s is:

An object is thrown vertically upward from the top of a 30 m high building with an initial speed of 20 m/s. The average velocity (in m/s) during the time interval  $t=0$  to  $t=5$  s is:

Select one:

- a. 13.8 downward
- b. 4.5 downward ✓ Well done!
- c. 0
- d. 13.8 upward
- e. 4.5 upward

The correct answer is: 4.5 downward

**Correct**

Marks for this submission: 2.0/2.0.

Two objects A and B are at the same height. A is projected vertically upwards with a speed of 20 m/s. At the same time B is projected vertically downward at 20 m/s. Which of the following statements is correct?

Select one:



Two objects A and B are at the same height. A is projected vertically upwards with a speed of 20 m/s. At the same time B is projected vertically downward at 20 m/s. Which of the following statements is correct?

Select one:

- a. A and B reach the ground with the same velocity. ✓ Well done!
- b. When reaching the ground B has higher velocity than A.
- c. A and B must have different velocities when reaching the ground.
- d. A and B reach the ground at the same time.
- e. A reaches the ground before B.

The correct answer is: A and B reach the ground with the same velocity.

Correct

Marks for this submission: 2.0/2.0.

The velocity of a particle moving along the x - axis is given by  $v(t) = 2t + 1$  where t is in seconds and v(t) in m/s. The average acceleration (in  $\text{m/s}^2$ ) over the time interval 0 to 2s is:

Correct

Marks for this submission: 2.0/2.0.

The velocity of a particle moving along the x - axis is given by  $v(t) = 2t + 1$  where  $t$  is in seconds and  $v(t)$  in m/s. The average acceleration (in  $\text{m/s}^2$ ) over the time interval 0 to 2s is:

Select one:

- a. 0
- b. -1.0
- c. 1.0
- d. 2.0 ✓ Well done!
- e. -2.0

The correct answer is: 2.0

Correct

Marks for this submission: 2.0/2.0.

An object is thrown vertically upwards with an initial speed of 30 m/s. After 4 s, the object is:

An object is thrown vertically upwards with an initial speed of 30 m/s. After 4 s, the object is:

Select one:

- a. moving up at 9.2 m/s
- b. moving up at 20 m/s
- c. moving down at 20 m/s
- d. moving down at 9.2 m/s ✓ Well done!
- e. at its maximum height

The correct answer is: moving down at 9.2 m/s

**Correct**

Marks for this submission: 2.0/2.0.

A car moving in one dimension travels from point A to point B at an average speed of 40 km/h. It then reverses direction and moves from point B back to point A at 20 km/h. Its average speed (in km/h) over the entire trip is:

Select one:



The correct answer is: moving down at 9.2 m/s

Correct

Marks for this submission: 2.0/2.0.

A car moving in one dimension travels from point A to point B at an average speed of 40 km/h. It then reverses direction and moves from point B back to point A at 20 km/h. Its average speed (in km/h) over the entire trip is:

Select one:

- a. 0
- b. 26.7 ✓ Well done!
- c. 40.0
- d. 60.0
- e. 20.0

The correct answer is: 26.7

Correct

Marks for this submission: 2.0/2.0.



The position of an object moving along the x-axis varies with time according to the equation  $x(t) = t^2 + 3t - 1$ . The average velocity (in m/s) of this object over the time interval 1 to 3 s is:

Select one:

- a. 7.0 ✓ Well done!
- b. -1.5
- c. 1.5
- d. -7.0
- e. 10

The correct answer is: 7.0

**Correct**

Marks for this submission: 2.0/2.0

[Finish review](#)

**Q1)** what is the pressure (in atm) at the base of a dam if the water is 200 m deep?

- A) 20.3                      B) 24.7                      C) 29.4                      D) 19.3                      E) 0

**Q2)** A supertanker filled with oil has a total mass of  $6 \times 10^8$  kg. If the dimensions of the ship are those of a box 300 m long, 80 m wide and 40 m high, determine how far (in m) the bottom of the ship is below sea water. (Assume density of sea water =  $1020 \text{ kg/m}^3$ )

- A) 10                      B) 15                      C) 18                      D) 19                      E) 25


**Q3)** Water flows (streamline, nonviscous) from point *a* to point *b* in the horizontal section shown in the figure. Which of the following statements is correct regarding the velocity *v*, pressure *P*, and flow rate at the two ends of the section?

- A)  $v_a < v_b$                       B)  $P_a > P_b$                       C)  $P_a < P_b$   
D)  $P_a = P_b$                       E)  $v_a = v_b$



**Q4)** A 4.0 cm radius horizontal pipe gradually narrows down to 2.0 cm. When water flows in this pipe the pressure in these two sections is 32.0 kPa and 24.0 kPa, respectively. What is the speed of the water through the smaller section?

- A) 4.1                      B) 1.0                      C) 3.5                      D) 17.0                      E) 5.4



**Q5)** A 2  $\mu\text{Ci}$  radioactive source emits neutrons each with 2.4 MeV energy. The radiated energy (in  $\mu\text{J}$  per hour) is: ( $\mu\text{Ci} = 10^{-6} \text{ Ci}$  and  $1 \text{ Ci} = 3.70 \times 10^{10} \text{ decays/sec}$ ,  $1 \text{ eV} = 1.6 \times 10^{-19} \text{ J}$ )

- A) 23                      B) 42                      C) 102                      D) 150                      E) 250

**Q6)** What is the activity (in decay/sec) of 8.8  $\mu\text{g}$  mass of  $^{124}\text{Cs}$ , which has a half-life of 30.8 s ?

- A)  $6.5 \times 10^{13}$                       B)  $9.6 \times 10^{14}$                       C)  $1.2 \times 10^{13}$                       D)  $8.8 \times 10^{13}$                       E)  $1.9 \times 10^{13}$

**Q7)** A radioactive isotope decays by  $\beta^-$  emission with a half-life of 1.0 day. The initial number of radioactive nuclei is 8000. The number of the remaining radioactive nuclei after 3 days is:

- A) 4000                      B) 1000                      C) 1500                      D) 2000                      E) 8000

**Q8)** A 2 mSv (milli sievert ) is equivalent to:

- A) 0.2 rem                      B) 0.2 rad                      C) 2 Gy                      D) 20 rem                      E) 20 rad


**Q9)** Three radioactive sources have the same activity. The decay modes, energy of each emitted particle and the corresponding RBE values are given in the following table. Arrange the three sources according to their danger to biological tissues from least to most dangerous. (Assume the **absorbed dose** from each source to be equal).

- A) 2, 3, 1  
B) 1, 2, 3  
C) 3, 2, 1  
D) 2, 1, 3  
E) 1, 3, 2

| Source | Decay mode      | Energy per emitted particle | RBE |
|--------|-----------------|-----------------------------|-----|
| 1      | protons         | 3MeV                        | 2   |
| 2      | Fast neutrons   | 1MeV                        | 10  |
| 3      | Alpha particles | 1 MeV                       | 20  |

**Q10)** A 70-kg laboratory technician exposed to  $\alpha$ -particles absorbs 0.03 mJ of energy. The relative biological effectiveness (RBE) for  $\alpha$ -particles is 20. What is his effective dose (in mrem)?

- A) 1.20                      B) 1.09                      C) 1.00                      D) 0.92                      E) 0.86



\* Which statement is correct?

- a) at  $\theta = 90^\circ \rightarrow T = m\vec{g}$
- b) at  $\theta = 90^\circ \rightarrow T = 0$
- c) at  $\theta = 0^\circ \rightarrow T = m\vec{g}$
- d) at  $\theta = 0^\circ \rightarrow T = 0$
- e) at  $\theta = 0^\circ \rightarrow$  The force that the wall exerts on the rod equals to zero

(III) Assume a liter of milk typically has an activity of 2000 pCi due to  $^{40}_{19}\text{K}$ . If a person drinks two glasses (0.5 L) per day, estimate the total effective dose (in Sv and in rem) received in a year. As a crude model, assume the milk stays in the stomach 12 hr and is then released. Assume also that roughly 10% of the 1.5 MeV released per decay is absorbed by the body. Compare your result to the normal allowed dose of 100 mrem per year. Make your estimate for (a) a 60-kg adult, and (b) a 6-kg baby.

1 L of milk  $\rightarrow 2000 * 3.7 * 10^{10} * 10^{-12}$  decay per s

\* a person drinks two glasses meaning (0.5 L) per day

\* milk stays in stomach for 12 h  $\rightarrow 43200$  s

\* to find the total energy we need to find number of decays  $\rightarrow$  time \* decay per s

$$\rightarrow 2000 * 3.7 * 10^{10} * 10^{-12} * 43200$$

$$= 3196800 \rightarrow \text{however this is for 1 L}$$

for 0.5 L we just multiply by 0.5

no. decays for one day  $\rightarrow 1598400$

now energy per decay is  $\rightarrow 1.5$  MeV

$$\begin{aligned} \text{tot energy per day} &\rightarrow 1598400 * 1.5 * 1.6 * 10^6 * 10^{-19} \\ &= 3.83616 * 10^{-7} \end{aligned}$$

the body only absorbs around 10% of the tot released energy  $\rightarrow 3.83616 * 10^{-7} \text{ J} * 0.1$

$$\rightarrow 3.83616 * 10^{-8} \text{ J per day}$$

now absorbed energy per year

$$\rightarrow 3.83616 * 10^{-8} * 365 = 1.4 * 10^{-5} \text{ J}$$

$$\text{dose for adult} = \frac{1.4 * 10^{-5}}{60} = 2.33 * 10^{-7} \text{ Sv/y}$$



// For child

$$= \frac{1.4 * 10^{-5}}{6} = 2.33 * 10^{-6} \text{ Sv/y}$$

$$6 \text{ or } 0.233 \text{ mrem/y}$$

$$\text{or } 2.33 * 10^{-2} \text{ mrem/y}$$

3 . There was a question about a child on a swing with a mass of 25 kg. He started from rest and his speed reached 2 m/s. Request the loss of mechanical energy due to friction..



$$\Delta ME = \Delta K + \Delta U$$

$$= \frac{v_f^2}{2} m - \frac{v_i^{\text{zero}}}{2} m + -mg(L - L \cos 30^\circ)$$

$L=2.4\text{m}$

**Question:** The speed of light in a medium is equal to 0.85 times the speed of light in water, given the index of refraction of water is 1.33. Calculate the index of refraction of this substance.

**Solution:**

1. **Speed of Light in Water:**

$$c_{\text{water}} = \frac{c}{n_{\text{water}}} = \frac{c}{1.33}$$

2. **Speed of Light in the Medium:**

$$c_{\text{medium}} = 0.85 \times c_{\text{water}} = 0.85 \times \frac{c}{1.33}$$

3. **Index of Refraction of the Medium:**

$$n_{\text{medium}} = \frac{c}{c_{\text{medium}}} = \frac{c}{0.85 \times \frac{c}{1.33}} = \frac{1.33}{0.85} \approx 1.56$$

**Final Answer:** The index of refraction of the substance is approximately 1.56.

**Question:** If the average pressure is  $1.33 \text{ N/cm}^2$  and the energy produced by the heart in 12 hours is sufficient to lift a body 15 m, with the mass of the body being 426 kg, what is the flow rate in  $\text{cm}^3/\text{s}$ ?

**Solution:**

1. Calculate the Work Done:

$$\text{Work} = \text{Mass} \times g \times \text{Height} = 426 \text{ kg} \times 9.81 \text{ m/s}^2 \times 15 \text{ m}$$

$$\text{Work} = 62607.9 \text{ J}$$

2. Convert 12 hours to seconds:

$$12 \text{ hours} = 12 \times 3600 \text{ s} = 43200 \text{ s}$$

3. Calculate Power:

$$\text{Power} = \frac{\text{Work}}{\text{Time}} = \frac{62607.9 \text{ J}}{43200 \text{ s}} \approx 1.45 \text{ W}$$

4. Calculate the flow rate: Using the pressure and power relationship:

$$\text{Flow rate} = \frac{\text{Power}}{\text{Pressure}} = \frac{1.45 \text{ W}}{1.33 \text{ N/cm}^2}$$

Converting pressure to  $\text{N/m}^2$  ( $1 \text{ N/cm}^2 = 10,000 \text{ N/m}^2$ ):

$$\text{Flow rate} = \frac{1.45}{1.33 \times 10^4} \approx 1.09 \times 10^{-4} \text{ m}^3/\text{s} = 109 \text{ cm}^3/\text{s}$$

**Final Answer:** The flow rate is approximately  $109 \text{ cm}^3/\text{s}$ .

**Question:** A body in the shape of the letter L has dimensions  $x_1 = 6$  and  $x_2 = 4$ . Calculate the center of mass (CM).

**Solution:** Assuming equal mass for both segments:

$$\text{CM} = \frac{m_1 x_1 + m_2 x_2}{m_1 + m_2}$$

Let  $m_1 = m$  and  $m_2 = m$ :

$$\text{CM} = \frac{m \cdot 6 + m \cdot 4}{m + m} = \frac{6 + 4}{2} = 5$$

**Final Answer:** The center of mass is at  $x = 5$  (in terms of  $m$ ).



**Question:** A cylinder is floating upright in water. The cylinder has a volume of 640 ml, a diameter of 7.6 cm, and a mass of 45 g when empty. The length of the cylinder above the water level is  $d$ . Determine  $d$  in cm.

**Solution:**

1. **Volume of Water in the Cylinder:** The cylinder is half full, so:

$$V_{\text{water}} = \frac{640 \text{ ml}}{2} = 320 \text{ ml} = 320 \text{ cm}^3$$

2. **Weight of Water:** The density of water is  $1000 \text{ kg/m}^3$  or  $1 \text{ g/cm}^3$ .

$$\text{Weight of Water} = 320 \text{ g} = 0.32 \text{ kg} \times 9.81 \text{ m/s}^2 = 3.136 \text{ N}$$

3. **Weight of the Cylinder:**

$$\text{Weight of Cylinder} = 45 \text{ g} = 0.045 \text{ kg} \times 9.81 \text{ m/s}^2 = 0.44145 \text{ N}$$

4. **Total Weight:**

$$\text{Total Weight} = 3.136 + 0.44145 \approx 3.57745 \text{ N}$$

5. **Buoyant Force:**

$$\text{Buoyant Force} = \text{Weight of Water displaced} = \text{Total Weight}$$

6. **Volume Displaced:**

$$V_{\text{displaced}} = \frac{3.57745 \text{ N}}{9.81 \text{ m/s}^2} \approx 0.364 \text{ kg} \rightarrow 364 \text{ cm}^3$$


7. **Total Volume of Cylinder:**

$$V_{\text{cylinder}} = \frac{640 \text{ ml}}{1000} = 640 \text{ cm}^3$$

8. **Length of Cylinder:** Volume of cylinder =  $\pi \left(\frac{d}{2}\right)^2 \cdot h$ . With  $d = 7.6 \text{ cm}$ :

$$h = \frac{640}{\pi \left(\frac{7.6}{2}\right)^2} \approx 11.82 \text{ cm}$$

8. Calculate the value of  $F$  so that the body begins to move, knowing that these forces are perpendicular to it



1. A  $7279 \text{ m}^3$  balloon is filled with hot air with a density of  $0.9447 \text{ kg/m}^3$ . The surrounding air has a density of  $1.25 \text{ kg/m}^3$ . How much weight (in N) can the balloon hold (including the balloon skin)?

- **Answer:** To find the weight the balloon can hold, we calculate the buoyant force, which is the difference between the density of the surrounding air and the density of the hot air inside the balloon. The buoyant force is given by:

$$\begin{aligned}\text{Weight} &= (\text{Density}_{\text{air, surrounding}} - \text{Density}_{\text{hot air}}) \times \text{Volume} \times g \\ &= (1.25 - 0.9447) \text{ kg/m}^3 \times 7279 \text{ m}^3 \times 9.81 \text{ m/s}^2 \\ &\approx 18431 \text{ N}\end{aligned}$$

2. The effective dose of alpha particles is  $0.14 \text{ mSv}$ . What is the absorbed dose? (RBE: 20)

- **Answer:** The absorbed dose can be calculated using the formula:

$$\begin{aligned}\text{Absorbed Dose} &= \frac{\text{Effective Dose}}{\text{RBE}} \\ &= \frac{0.14 \text{ mSv}}{20} \\ &= 0.007 \text{ mSv} = 7 \text{ mGy}\end{aligned}$$

3. An iodine isotope has a half-life of 8.04 days. A sample for treatment had an activity of 5 mCi before shipment, and when received, it had an activity of 2.1 mCi. What is the time interval (in days) between shipment and receipt?

- **Answer:** Using the decay formula:

$$\text{Activity} = \text{Activity}_0 \times e^{-\lambda t}$$

where  $\lambda$  is the decay constant:

$$\lambda = \frac{\ln(2)}{T_{1/2}}$$

Rearranging to solve for  $t$ :

$$t = \frac{\ln\left(\frac{\text{Activity}_0}{\text{Activity}}\right)}{\lambda}$$

$$= \frac{\ln\left(\frac{5}{2.1}\right)}{\frac{\ln(2)}{8.04}}$$

$$\approx 101 \text{ days}$$

4. A diverging lens with focal length  $f$  has an object placed at  $3f$  from the lens. What is  $d_i$  (the image distance)?

- **Answer:** For a diverging lens, the lens equation is:

$$\frac{1}{f} = \frac{1}{d_o} + \frac{1}{d_i}$$

where  $d_o = 3f$ :

$$\frac{1}{f} = \frac{1}{3f} + \frac{1}{d_i}$$

Solving for  $d_i$ :

$$\frac{1}{d_i} = \frac{1}{f} - \frac{1}{3f} = \frac{2}{3f}$$

$$d_i = -\frac{3f}{2}$$

$$= -\frac{3}{4}f$$

5. A water tank open to the atmosphere has a hole 21 m below the water level. The hole's area is  $10^{-4} \text{ m}^2$ . What is the volume flow rate?

- Answer: Using Torricelli's theorem:

$$v = \sqrt{2gh}$$

where  $h = 21 \text{ m}$  and  $g = 9.81 \text{ m/s}^2$ :

$$v = \sqrt{2 \times 9.81 \times 21} \approx 20.5 \text{ m/s}$$

The volume flow rate  $Q$  is:

$$\begin{aligned} Q &= A \times v = 10^{-4} \text{ m}^2 \times 20.5 \text{ m/s} \approx 2.05 \times 10^{-3} \text{ m}^3/\text{s} \\ &\approx 1.5 \times 10^{-3} \text{ m}^3/\text{s} \end{aligned}$$

6. A converging lens with a power of 10 diopters. If the object distance  $d_o$  is 8 cm, what is  $d_i$  (the image distance)?

- Answer: The power  $P$  is related to the focal length  $f$  by:

$$P = \frac{1}{f}$$

Thus,  $f = \frac{1}{10} \text{ m} = 0.1 \text{ m} = 10 \text{ cm}$ . Using the lens equation:

$$\frac{1}{f} = \frac{1}{d_o} + \frac{1}{d_i}$$

$$\frac{1}{10} = \frac{1}{8} + \frac{1}{d_i}$$

Solving for  $d_i$ :

$$\frac{1}{d_i} = \frac{1}{10} - \frac{1}{8} = \frac{-1}{40}$$

$$d_i = -40 \text{ cm}$$

7. A box with a mass of 3 kg is on a 35-degree incline with a static friction coefficient of 0.4. What is the minimum applied force perpendicular to the incline needed to prevent the box from sliding?

- **Answer:** To prevent sliding, the applied force must counteract the component of gravitational force parallel to the incline. The normal force  $N$  on the incline is:

$$N = mg \cos \theta$$

The frictional force  $F_f$  is:

$$F_f = \mu N = \mu mg \cos \theta$$

The component of gravitational force parallel to the incline is:

$$F_{\text{parallel}} = mg \sin \theta$$

Setting  $F_f = F_{\text{parallel}}$ :

$$\mu mg \cos \theta = mg \sin \theta$$

Solving for the applied force perpendicular to the incline:

$$F_{\text{applied}} = F_{\text{parallel}} - \mu N$$

$$F_{\text{applied}} = mg(\sin \theta - \mu \cos \theta)$$

10. For a material with  $R = 6 \times 10^{11}$  and  $N = 2 \times 10^{15}$  atoms, what is the half-life (in minutes)?

- **Answer:** The half-life  $T_{1/2}$  can be calculated using:

$$T_{1/2} = \frac{\ln 2}{\lambda}$$

where  $\lambda = \frac{R}{N}$ :

$$\lambda = \frac{6 \times 10^{11}}{2 \times 10^{15}} \approx 3 \times 10^{-4} \text{ s}^{-1}$$

$$T_{1/2} = \frac{\ln 2}{3 \times 10^{-4}} \approx 2315 \text{ s} \approx 38.5 \text{ min}$$

11. A question involving the application of Torricelli's theorem to find the volume flow rate: The equation  $v = \sqrt{2gh}$  must be used to find the velocity of the liquid exiting the small opening. Then the velocity must be multiplied by the cross-sectional area (given) to find the volume flow rate.

- **Answer:** Apply Torricelli's theorem:

$$v = \sqrt{2gh}$$

The volume flow rate  $Q$  is:

$$Q = A \times v$$



12. The speed of light in a medium is 82% of the speed of light in water. Given that the refractive index of water is 1.33, what is the refractive index of that medium?

- **Answer:** The refractive index  $n$  is given by:

$$n = \frac{c_{\text{water}}}{c_{\text{medium}}}$$

where  $c_{\text{medium}} = 0.82 \times c_{\text{water}}$ :

$$n = \frac{1.33}{0.82} \approx 1.62$$

13. A converging lens with a power of 1 diopter. If the object was placed 30 cm away from the lens, how far was the image from the lens?

- **Answer:** The power  $P$  is related to the focal length  $f$  by:

$$P = \frac{1}{f}$$

Thus,  $f = \frac{1}{1} \text{ m} = 1 \text{ m} = 100 \text{ cm}$ . Using the lens equation:


$$\frac{1}{f} = \frac{1}{d_o} + \frac{1}{d_i}$$
$$\frac{1}{100} = \frac{1}{30} + \frac{1}{d_i}$$

Solving for  $d_i$ :

$$\frac{1}{d_i} = \frac{1}{100} - \frac{1}{30} = \frac{-7}{300}$$
$$d_i \approx -40 \text{ cm}$$

Q14

Q) Ray of light passes from Medium X into the liver.  
The speed of light in the liver gets reduced by 10%.  
What is the depth of the liver (d)?




Q15

-What is horsepower?


Ans: the rate which an object can do work

Q16



the weight of the load  
is 1400 N and a  
object weight 2000 N  
what is the magnitude  
of tension force?

Q17



what is the  
magnitude of  $(d)$ ?