



Carbohydrates Metabolism




Review of Carbohydrates
Digestion and absorption of carbohydrates

Dr. Diala Abu-Hassan




Sugars are either
aldoses or ketoses

A Aldehyde group



Ribose
Glucose


B Keto group



Ribulose
Fructose

Examples of monosaccharides found in humans

Enantiomers



Generic names

3 carbons: trioses

4 carbons: tetroses

5 carbons: pentoses

6 carbons: hexoses

7 carbons: heptoses

9 carbons: nonoses

Examples

Glyceraldehyde

Erythrose

Ribose

Glucose

Sedoheptulose


Neuraminic acid

Sugars have Isomers


Epimers are isomers:

Changing the orientation of one hydroxyl group will produce a different sugar

Glucose and Fructose are isomers




Alpha and Beta Sugars (Anomers)




Disaccharides


Sugars made of two monosaccharide units joined by a glycosidic bond




Maltose: a disaccharide made from two glucose units




Glycosidic bond is cleaved by glycosidase enzyme





Digestion of Carbohydrates

Starch Digestion



Mucosal cell membrane-bound enzymes

ENZYME	Bond Cleaved	Substrates
Isomaltase	α 1 $\rightarrow$ 6	Isomaltose
Maltase	α 1 $\rightarrow$ 4	Maltose
Sucrase	α 1 $\rightarrow$ 2	Sucrose
Lactase	β 1 $\rightarrow$ 4	Lactose
Trehalase	α 1 $\rightarrow$ 1	Trehalose
Exoglycosidase (Glucoamylase)	α 1 $\rightarrow$ 4 and α 1 $\rightarrow$ 6	Starch

Sucrase-isomaltase complex and Glucoamylase

* Sucrase + isomaltase

Single protein → complex of two
associated subunits

- Sucrase-maltase

- Isomaltase-maltase

Together 80% of the maltase activity

* Maltase + exoglycosidase
(glucoamylase): no split




FIG. 27.5. The major portion of the sucrase-isomaltase complex, containing the catalytic sites, protrudes from the absorptive cells into the lumen of the intestine. Other domains of the protein form a connecting segment (stalk) and an anchoring segment that extends through the membrane into the cell. The complex is synthesized as a single polypeptide chain that is split into its two enzyme subunits extracellularly. Each subunit is a domain with a catalytic site (distinct sucrase-maltase and isomaltase-maltase sites). In spite of their maltase activity, these catalytic sites are often called just *sucrase* and *isomaltase*.


Clinical Hint: Abnormal Degradation of disaccharides

1. Sucrase-isomaltase deficiency:


Causes:

- Genetics
- Variety of intestinal diseases
- Malnutrition
- Injury of mucosa i.e by drugs
- Severe diarrhea

c SI carbohydrate breakdown




CSID and IBS risk variants



Clinical Hint: Abnormal Degradation of disaccharides

2. Lactase deficiency: ½ world's population

- ✓ Lactase reached maximal activity @ 1 month of age
- ✓ Declines ----- >> adult level at 5 to 7 year of age
- ✓ 10 % of infant level
- ✓ 1 cup of milk (9 grams of lactose) → loss of 1 liter of extracellular fluid



Absorption of Sugars

- ✓ Polar molecules can not diffuse
- ✓ Na^+ -independent facilitated diffusion transport
- ✓ GLUT 1-----GLUT 14
- ✓ Glc. Movement follows concentration gradient
- ✓ Two conformational states
- ✓ **Glucose transporter inhibitors** are used to decrease sugar transport detected in many types of cancer because the expression of these transporters is upregulated in cancer.





Table 27.5 Properties of the GLUT 1 to GLUT 5 Isoforms of the Glucose Transport Proteins

<i>Transporter</i>	<i>Tissue Distribution</i>	<i>Comments</i>
GLUT 1	Human erythrocyte Blood–brain barrier Blood–retinal barrier Blood–placental barrier Blood–testis barrier	Expressed in cell types with barrier functions; a high-affinity glucose transport system
GLUT 2 Glucose, galactose and fructose	Liver Kidney Pancreatic β -cell Serosal surface of intestinal mucosa cells (Basolateral surface)	A high-capacity, low-affinity transporter May be used as the glucose sensor in the pancreas
GLUT 3	Brain (neurons)	Major transporter in the central nervous system, a high-affinity system
GLUT 4	Adipose tissue Skeletal muscle Heart muscle	Insulin-sensitive transporter. In the presence of insulin, the number of GLUT 4 transporters increases on the cell surface; a high-affinity system
GLUT 5 Fructose	Intestinal epithelium Spermatozoa	This is actually a fructose transporter Na independent
GLUT 7	Glucogenic tissues	at endoplasmic reticulum membrane




Insulin stimulates transport of glucose into muscle and adipose tissues

Na⁺ monosaccharide cotransporter system (SGLT)

- Against concentration gradient (requires energy).

* Small intestine:
Active uptake from
lumen of intestine.

- Kidney:
reabsorption of
glucose in proximal
tubule.
- For glucose and
galactose
absorption



, Na⁺-glucose cotransporters



, Facilitated glucose transporters



, Na⁺,K⁺-ATPase

- SGLT inhibitors (**gliflozins or flozins**), are a class of antidiabetic agents that promote renal glucose excretion independently of insulin.