

Lecture 3

Bacterial Growth& physiology





Bacterial Growth

Definitions

B. Reproduction

culture media


A) Definition

B) Classification

C) Types

Bacterial growth curve


Bacterial Growth: Definition




I) Bacterial Growth

Indicated by

a) Turbidity of the fluid media







I) Bacterial Growth

Indicated by

b) Colonies on solid media

(Macroscopic product)





2) Colony (Macroscopic product)

Single bacterium

On solid media

After 20-30 division

Binary fission





So the appearance of colony will be after 20-30 division of single bacterium


2) Colony

After 20-30 division

Binary fission


I Million

(2^20)




3) Generation time (doubling time)

13min (V.cholerae)



Binary fission









Separation of 2 strands


(ssDNA attached to mesosome)




2 Separation of 2 strands



3 Separate ssDNA & become dsDNA




Formation of division septum





Cell separation






Bacterial culture media

Bacteria grow

(In vitro)



Need nutrients for growth



Artificial








Isolation & diagnosis (Causative agent)




Prepare vaccine & Other product



For Selection

proper antibiotics




Classification of media

Liquid



Solid



Types of media

- 1) Simple media
- 2) Enriched media
- 3) Selective media
- 4) Differential media



Basic requirement for growth of most bacteria



A) Peptone water

Peptone + 0.5% NaCl

Enhancement

Sugar media



B) Nutrient broth

Meat extract

Enhancement



C) Nutrient agar plate

Nutrient broth + 2%

agar agar

(Seaweed)



Staph. aureus

Fastidious bacteria
Need blood, serum for
growth



A) Blood agar

Nutrient agar heated at

45°C

(semisolid)


+ sheep blood




A) Blood agar

Streptococci

Strept. pyogenes






Hemolysis on blood agar:

- A. Complete (beta) hemolysis:
 - Staphylococcus aureus
 - Streptococcus pyogenes

- C. No (gamma) hemolysis:
- Enterococci.

- B. Partial (alpha) hemolysis:
- Streptococcus viridans
- pneumococci.




B) Chocolate agar

Nutrient agar heated at 100°C, add blood

Hb Heat Haematin

(Chocolate)






B) Chocolate agar

Haemophilus Neisseria






Selective media

Allow a certain organism to grow
(Selective) &
inhibits the growth of others






Selective media

Lowenstein Jensen medium

Malachite green

Mycobacterium tuberculosis






Selective media

Blood tellurite agar

Potassium tellurite

C.diphtheriae






Differential media

Selective + Indicator

Allow a certain organism to grow

Indicator to differentiate (change in visibly)



Purpose Purpose

MacConkey's agar

Bile (Enterobacteria)

Lactose= test sugar

Peptone

Neutral red =pH indicator




Pink

Pale

Differential media

Mannital salt agar
(high salt 7.5% NaCl)
Phenal red =pH indicator



Staphylócoccus aureus

Differential media

Thiosulfate-Citrate-Bile-Sucrose Agar. (TCBS)


Thiosulphate

Bile

Citrate

Sucrose as test sugar

Bromothymol blue




- I) Lag phase
- 2) Log phase
- 3) Stationary phase
- 4) Decline phase

If a small number


of bacteria are inoculated

into a liquid nutrient

medium



72h



Lag phase

No. of bacteria




Lag phase

No. constant

Log phase

No. of bacteria



Stationary phase

No. of bacteria

3

Stationary phase


Constant number


No. of division = No. of

death

(Waste product)

Decline phase





Bacterial growth requirements

Growth Requirements

- A) Nutrition
- B) Gaseous
- C) Temp. & pH

A) Nutrition

Maintenance of bacterial growth



A) Nutrition

I- Autotrophic

auto = self

Trophic=nutrition

2- Heterotrophic

hetero = different


Trophic = nutrition


Autotrophic

CO₂

Ammonium

Utilize simple inorganic substance






Complex organic materials

(Saprophytic)

Autotrophic

No medical importance



Heterotrophic

These bacteria require complex preformed organic substances e.g. sugars, proteins etc.

Parasitic

Medical important

Living host


Growth Requirements

Gaseous requirements

O₂ requirement, bacteria are classified into 5 groups

02 requirement

- I) Obligate aerobes
- 2) Obligate anaerobes
- 3) Facultative anaerobes
- 4) Micro-aerophilic
- 5) Aero-tolerant



Respiration

Glucose catabolism

Energy production

Aerobic respiration

 (0_2)

Anaerobic respiration

 (NOO_2)



I- Obligate aerobes (Aerobic respiration)

Presence of 0₂

Absence of O₂


Growth

No growth


e.g. Pseudomonas aeruginosa

I- Obligate aerobes (Aerobic respiration)

Aerobic respiration



For production Energy (ATP)



Glucose catabolism (glycolysis)

I- Obligate aerobes (Aerobic respiration)

Oxidative phosphorylation

$$0_2 \longrightarrow Carry 2H+ \longrightarrow H_2 0+34 ATP$$


I-Obligate aerobes

Highly toxic molecules

Superoxide (0₂)

Superoxide dismutase

(H₂ O₂)



Catalase

2- Obligate anaerobes


Presence of 02

Absence of 02

No growth

Growth

Bacteroides fragilis



Lack Superoxide dismutase & Catalase

The organism used inorganic molecules




Carry H+


13 ATP + 4 ATP 17 ATP

Lack

Superoxide dismutase

Catalase







3-Facultative anaerobes

Presence of O₂






Rate of growth



Most bacteria

3-Facultative anaerobes




4-Micro-aerophilic

Presence of O₂

superoxide & H₂ O₂

No growth

Low O₂




Low superoxide dismutase & catalase

Campylobacter
Helicobacter




Low 02



Superoxide dismutase



Absence of 02



Growth

Cl.perfringens

Growth Requirement: CO2 requirements

 CO_2 (0.03%)

Present in air is sufficient

CO₂ (5-10%)

(Capnophilic)


Neisseria

Brucella

Growth Requirement: Hydrogen ion (pH)

pH(7.2 - 7.4)

(Most bacteria)




Hydrogen ion (pH)

Alkaline (pH 9)



Vibrio cholerae

Acidic (pH 4)



Lactobacilli

Growth Requirement: Temperature

Mesophilic

(20-45)

(Most bacteria)

Psychrophilic

(0-15)

Thermophilic

(55-65)