
بسم الله الرحمن الرحيم (وَفَوْقَ كُلِّ ذِي عِلْمٍ عَلِيمٌ)



Microbiology | Lecture #5

Bacterial Structure pt.2



Written by: Dopamine 023


Reviewed by: NST

Objectives

Structures outside the cell wall

- 1) Capsule
- 2) Flagella
- 3) Pili
- 4) Spore formation

Capsule - Definition




A capsule is a wall made of carbohydrates that surrounds the cell wall.

Capsule - Definition

The capsule is an extra layer since it is NOT present in ALL bacteria.

Gelatinous (Viscous) layer covering cell wall of some bacteria



Extra layer


Usually Polysaccharides

EXCEPT



Polypeptides

(B. anthracis)



Variation of Capsule



Caused by different

(Arrangement of

Polysaccharides)


Sucrose	Mannose	Lactose
Mannose	Sucrose	Mannose
Lactose	Sucrose	Mannose

e.g. 91 types of

Str. pneumoniae

Due to different polysaccharides arrangement

Do Not stained by Gram stain






That is why we see Unstained halo around the

organism

This is one of the mechanisms used to identify bacteria that has capsules.

Quellung reaction (swelling)



Capsule

The name differs depending on the binding of the substance to the cell wall.


Capsule

Glycocalyx

Has fibril extensions help it adhere

Slime layer

These components are similar in their loose, unorganized attachment




If the substance is highly attached to the cell wall, we call it a capsule

Capsule

Tightly, organized bound around all cell wall


Firmly adherence to surface organism



Glycocalyx

(Slime layer)

Loosely & unorganized attached



Glycocalyx


Fibrils extending

Makeit

It adhere firmly to skin, heart, etc

e.g. Strept. mutans

It adheres to the host cell NOT the bacterial capsule



Loosely & unorganized attached




Protect Cell wall

From

Bacteriophage

Infective virus that has specific receptors on the bacterial wall. When the cell wall is surrounded by capsules, it prevents bacteriophages from binding to the bacteria resulting in no infection.








Protect Cell wall From Complements (in immunology)

(e.g lectin &alternative pathway)
The complement system must adhere to parts of the bacterial cell wall to start working. The capsule prevents the complement from binding resulting in more infection.








Protect Cell wall

From Lysozymes

= Enzymes that degrade bacterial cell wall.
Presence of the capsule prevents the breakdown of the cell wall by lysozymes.





Virulence factor: the ability of a pathogen to cause the disease.




Prevent phagocytosis


Capsule is a virulence factor


(Virulence)

The capsule protects the cell wall from phagocytosis to protect itself. (by "running away" from the phagocyte)



Phagocytosis Prevention by the bacteria







Capsules are formed

in VIVO ONLY

When the bacteria enter the host cell, they start forming the capsule by producing the components inside and secreting them to the outer surface of the bacterial cell wall.







The sugars in the bacteria undergo fermentation and end up releasing acids that result in formation of dental caries.


Attachment

(Glycocalyx)

تسوس الأسنان

Dental caries

The fibril extensions bind to any medical device (like implants, prosthetics..) in this case they adhere to the tooth enamel





Attachment


(Glycocalyx)

This is a virulence factor since the glycocalyx high adherence (in this case to the heart) leads to diseases.



Prosthetic heart

valves





Development of

vaccine

This is performed by extracting the capsule of "Haemophilus Influenzae b" bacteria and binding it to a protein.




Long thick threads like (filamentous),

formed from protein (flagellin)

(HAg)

Every flagella present in any bacteria is symbolized by (H Ag)




Seen by EM


(20nm)

Very small in size




Polar






Peri/trichous



The main function of the flagella is

The organ of motility



Motility

Gram -ve bacteria have HIGHER motility than Gram +ve bacteria because they have more rings

Gram -ve bacteria have 4 rings Gram +ve bacteria have 2 rings

Basal

body

Ring

For the filament to move, the rings must move, this causes the flagella to move as well as the bacteria.

Proton motive force

Filament

₽Hook

Outer

membrane

Peptidoglycan

Cell membrane

STRUCTURE OF FLAGELLA

This force makes the ring move

Tactic response (Taxis)

(Stimulus)


(movement of bacteria to

toward (+ve) or away (-ve)

from stimulating agent)

Positive chemotactic response ->toward material Negative chemotactic response ->away from material.

This response is due the the cell's chemotactic system where cell membrane send signals to direct flagella toward beneficial materials and away from harmful ones.



Tactic response (Taxis)

Stimulating agent


Two types of tactic response

Chemo Taxis

If the stimulus is chemical

Photo Taxis

If the stimulus is light



Chemical


Light

Axial Filaments

Endoflagella

In spirochetes (Example)


- > Some bacteria posses internal flagella instead of outside the bacteria.
- > This is known as endoflagella also referred as axial filaments.
- > These bacteria has wave like movements.



Pili (Fimbriae)

Short and thin
Hair like formed from
protein

(Pilin) Name of protein



Pili

only Seen by EM



Pili


Two types of pili:

A) Ordinary pili

(Attachment)


B) Sex pili

(Genetic transfer)




Because this pili act as defending weapon and help in adhering of bacteria to the host cell in the first step of infection (establishing a disease).

Ordinary Pili




Ordinary Pili

This photo from urethra



Bacteria can resist the fleshing of the urine.



Vegetative bacteria: bacteria that divide and replicate INSIDE THE HOST CEIL







specific disinfectant, etc.

(Outside)

Outside the host cell





Forming highly resistant resting

phase (Endospores) in VITRO

Only 2 types of bacteria can form spores:


OUTSIDE THE HOST CELL

Bacillus

Clostridium

Resting phase:

Means in dormant situation (do not perform any divisions or reproduction)




Occur to unfavorable conditions e.g.

High temp.


Drying

Depletion of nutrition



Formed outside the body (in VITRO)


Can not stained by ordinary stain Instead they have specific stains




Spores are:

Highly resistant to dryness,


heat & Disinfectant





It can remains for a long time until it finds the host

After entering the host





Ca+2 &
Diplconic acid


Explained in next slide

Multiple membranes


The Process of Spore Formation







Formation of Endospore




The Process of Spore Formation

- 1) First DNA replication occur, where each copy of the DNA moves to opposite sides of the bacterial cell.
- 2) Then multiple layers of the cell membrane and peptidoglycan form, producing calcium and dipicolinic acid, both contributing to the tough protective layers. (

 طبقة جيرية قاسية).
- 3) Next, a cortex forms, followed by the development of a spore coat, which contains over 80 types of proteins. Finally, an exosporium forms around the spore which is a collagen like glycoprotein. Once the spore is fully developed, the bacterium can exit the host cell and remain dormant, potentially surviving for centuries in harsh conditions.

Layers that forms from inside to outside:

Cortex → **Spore Coat** → **Exosporium**



Madefrom >80 proteins.


Germination التبرعم




Germination of spores occurs when the bacteria find suitable conditions like availability of water.

In germination, bacteria will break down all the layers (cortex, spore coat, exosporium) and return to become vegetative bacteria and begin their activities inside the host cell.

The Process of Germination





Position of spores

Depends on the location of spores.




B. anthracis

Central & Oval



Cl. perfringens

Sub-terminal & Oval



Cl. Tetani

Terminal & Spherical

رسالة من الفريق العلمي:

دعاء سيد الاستغفار

اللَّهُمَّ أَنْتَ رَبِّي لا إِلَهَ إلا أَنْتَ، خَلَقْتَنِي وَأَنَا عَبْدُكَ، وَأَنَا عَلَى عَهْدِكَ وَوَعْدِكَ مَا اسْتَطَعْتُ، أَعُوذُ بِكَ مِنْ شَرِّ مَا صَنَعْتُ، أَبُوءُ لَكَ بِنِعْمَتِكَ عَلَى، وَأَبُوءُ لَكَ بِذَنْبِي، فَاغْفِرْ لِي؛ فَإِنَّهُ لا يَغْفِرُ الذُّنُوبَ إلا أَنْتَ.

For any feedback, scan the code or click on it.



Corrections from previous versions:

Versions	Slide # and Place of Error	Before Correction	After Correction
V0 → V1			
V1 → V2			