

# Introduction to Microbiology

---

Anas Abu-Humaidan  
M.D. Ph.D.



# Overview

Bacterial genera that will be discussed this lecture are curved, gram-negative rods. that usually infect the GI tract:

- *Campylobacter*
- *Helicobacter*
- *Vibrio*

We also briefly discuss anaerobic gram negative rods


- *Bacteroids*

We will also discuss bacteria that do not have a cell wall

- *Mycoplasma*


# *Campylobacter*

- Small (0.2 to 0.5  $\mu\text{m}$  wide and 0.5 to 5.0  $\mu\text{m}$  long), motile, **curved, gram-negative rods**
- *Campylobacter* is the most common cause of **bacterial gastroenteritis**, with ***Campylobacter jejuni*** responsible for most infections.
- The organisms grow best in an atmosphere of reduced oxygen (5% to 7%) and increased carbon dioxide (5% to 10%) these properties are referred to as **microaerophilic**. ***C. jejuni* grows better at 42° C than at 37° C.**
- Express **lipooligosaccharides** (LOSs lack O-antigen in LPS)
- The organisms are killed when exposed to gastric acids, so conditions that decrease or neutralize gastric acid secretion favor disease
- *C. jejuni* GI disease characteristically produces **histologic damage to the mucosal surfaces of the jejunum** and other parts of the intestine



**FIGURE 28-1** Mixed culture of bacteria from a fecal specimen. *Campylobacter jejuni* is the thin, curved, gram-negative bacteria (arrow).

obligate  
aerobes




A

obligate  
anaerobes




B

facultative  
anaerobes




C

aerotolerant  
anaerobes



D

microaerophiles



E

Figure 2. Diagram of bacterial cell distribution in thioglycolate tubes.

# *Campylobacter*

- *Campylobacter* infections are **zoonotic**, with a variety of animals serving as reservoirs. **Contaminated poultry** are responsible for more than half of the *Campylobacter* infections in developed countries.
- **Uncommon for the disease to be transmitted by food handlers.**
- Present most commonly as **acute enteritis** with **diarrhea** (stools may be bloody on gross examination), **fever**, and **severe** abdominal pain.
- **Guillain-Barré syndrome** and **reactive arthritis** are well-recognized complications of *Campylobacter* infections (although uncommon). Probably through molecular mimicry.
- A **presumptive identification** of isolates is based on **growth under selective conditions**, typical **microscopic morphology**, and positive oxidase and catalase tests.

Microbiological finding	No. (%)	95% CI, %
Stool pathogen isolated <sup>a</sup>	168 (30.6)	27–35
<i>Shigella</i> species	84 (15.3)	12–19
<i>Salmonella</i> species	32 (5.8)	4–8
<i>Campylobacter</i> species	34 (6.2)	4–8
STEC	14 (2.6) <sup>b</sup>	1–4
Other enteropathogens <sup>c</sup>	9 (1.6)	1–3

**NOTE.** STEC, Shiga toxin-producing *Escherichia coli*.

<sup>a</sup> Three patients' stool specimens yielded 2 enteropathogens; each *Shigella* plus 1 each *Plesiomonas* or *Salmonella* species or *E. coli* O111.

<sup>b</sup> Includes 6 confirmed and 8 possible STEC cases.


<sup>c</sup> *Vibrio* (4), *Yersinia* (4), *Plesiomonas* (1) species.


Microbiological findings among US emergency department patients presenting with 549 episodes of bloody diarrhea at 11 *EMERGE*nc ID NET sites.

# *Helicobacter*

- **spiral gram-negative rods** resembling campylobacters, All gastric helicobacters, including *H. pylori*, are highly **motile** (corkscrew motility) and produce an abundance of **urease**
- Growth of *H. pylori* and other helicobacters requires a **complex medium** in **microaerophilic** conditions. *H. pylori* adheres to gastric mucosa and is usually not recovered in stool or blood specimens
- *H pylori* use their motility, chemotaxis, urease production, and other mechanisms to **adapt to the acidic conditions of the stomach** and colonize a narrow protected niche near the surface of epithelial cells
- **Humans are the primary reservoir for *H. pylori***, and colonization is believed to persist for life unless the host is specifically treated. Transmission is most likely via the **fecal-oral route**.

## The structure of *Helicobacter pylori*






Localized tissue damage is mediated by urease byproducts, **mucinase**, **phospholipases**, and the activity of **vacuolating cytotoxin A (VacA)**, a protein that after penetration into epithelial cells damages the cells by producing vacuoles. **cytotoxin-associated gene (cagA)** interferes with the normal cytoskeletal structure of the epithelial cells

# *Helicobacter*


- Colonization with *H. pylori* invariably leads to **gastritis**
- The **acute phase** of gastritis is characterized by a **feeling of fullness, nausea, vomiting, and hypochlorhydria**.
- Can evolve into **chronic gastritis**, with disease confined to the gastric antrum or involve the entire stomach
- Chronic gastritis will progress to develop peptic ulcers. The ulcers develop at the sites of intense inflammation, commonly involving the junction between the corpus and antrum (**gastric ulcer**) or the proximal duodenum (**duodenal ulcer**).
- *H. pylori* is responsible for 85% of the gastric ulcers and 95% of the duodenal ulcers.
- Chronic gastritis increases the risk of **gastric cancer and MALT lymphoma** (mucosa-associated lymphoid tissue B-cell lymphomas).



# *Helicobacter*

- Since *H. pylori* adheres to gastric mucosa, *H. pylori* can be detected by **histologic examination of gastric biopsy** specimens, but identification is usually done by non-invasive methods. A number of **polyclonal and monoclonal immunoassays for *H. pylori* antigens excreted in stool** have been developed and demonstrated to have sensitivities and specificities exceeding 95%.

Serology	Widely available Least expensive of available tests	Positive results may reflect previous rather than current infection Not recommended for confirming eradication
Urea breath test	High negative and positive predictive values Useful before and after treatment	False-negative results possible in the presence of proton pump inhibitors or with recent use of antibiotics or bismuth preparations Considerable resources and personnel required to perform test
Stool antigen test	High negative and positive predictive values with monoclonal antibody test Useful before and after treatment	Process of stool collection may be distasteful to patient False-negative results possible in the presence of proton-pump inhibitors or with recent use of antibiotics or bismuth preparation





# THE LANCET


Volume 321, Issue 8336, 4 June 1983, Pages 1273-1275

## Letters to the Editor

### UNIDENTIFIED CURVED BACILLI ON GASTRIC EPITHELIUM IN ACTIVE CHRONIC GASTRITIS

J Robin Warren <sup>a</sup>, Barry Marshall <sup>b</sup>



#### The structure of *Helicobacter pylori*




In 2005, the Karolinska Institute in Stockholm awarded the Nobel Prize in Physiology or Medicine to Marshall and Robin Warren, his long-time collaborator, "for their discovery of the bacterium *Helicobacter pylori* and its role in gastritis and peptic ulcer disease"

# *Vibrio*

- **Gram-negative, facultatively anaerobic, fermentative rods**, characterized by a **positive oxidase reaction** and the presence of **polar flagella** .
- *Vibrio* species can grow on a variety of simple media within a broad temperature range (from 14° C to 40° C). And tolerate a wide range of pH (e.g., pH of 6.5 to 9.0) but are **susceptible to stomach acids**.
- All species of *Vibrio* **require sodium chloride (NaCl)** for growth. Most species are halophilic (“salt-loving”).
- *Vibrio* species, including *V. cholerae*, grow naturally in **estuarine and marine environments** worldwide.
- Pathogenic vibrios can also flourish in waters with chitinous **shellfish**



### Environmental



(a)

### Pathogenic

Ingestion of contaminated food or water


Biofilm protects against gastric acid shock

Acid-sensitive cells conserve energy. Many cells die

Hyperinfectious biofilm and planktonic cells shed

Late infection

Early infection



(b)

# *Vibrio*

- All strains possess **lipopolysaccharides** consisting of lipid A (endotoxin), core polysaccharide, and an **O polysaccharide** side chain.
- The O polysaccharide is used to subdivide *Vibrio* species into **serogroups**, ***V. cholerae* O1 and O139** produce **cholera toxin** and are associated with epidemics of cholera. Other strains of *V. cholerae* generally do not produce cholera toxin and do not cause epidemic disease.
- Cholera is spread by **contaminated water and food** rather than direct person-to-person spread, because a **high inoculum** (e.g.,  $>10^8$  organisms) is required to establish infection in a person with normal gastric acidity.
- Cholera is usually seen in communities with **poor sanitation**. **Immunoassays** for the detection of cholera toxin or the O1 and O139 lipopolysaccharides are used for the diagnosis of cholera in endemic areas.
- It is estimated that **3 to 5 million cases** of cholera and 120,000 deaths occur worldwide **each year**. Seven major pandemics of cholera have occurred since 1817, resulting in thousands of deaths and major socioeconomic changes.

## Countries reporting cholera, 2010–2015



The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

Data Source: World Health Organization  
Map Production: Information Evidence  
and Research (IER)  
World Health Organization



© WHO 2016. All rights reserved.



Map of London, 1854. Water-distribution systems, which John Snow investigated comparing cholera cases among consumers of water of two suppliers depending on the site of their water intake from the Thames River.

# Vibrio


- The majority of individuals exposed to toxigenic ***V. cholerae* O1** have asymptomatic infections or self-limited diarrhea; however, some individuals develop severe, rapidly fatal diarrhea. *V. cholerae* O1 does not produce a capsule, so infections with this organism do not spread beyond the confines of the intestine.
- The clinical manifestations of cholera begin an average of 2 to 3 days after ingestion of the bacteria (can be <12 hours), with the abrupt onset of watery diarrhea and vomiting. Fever is rare.
- The feces-streaked stool specimens become colorless and odorless, free of protein, and speckled with mucus (**“rice-water” stools**).
- Patients with cholera must be promptly treated with **fluid and electrolyte replacement** before the resultant massive fluid loss leads to hypovolemic shock.

Species	Virulence Factor	Biological Effect
<i>V. cholerae</i>	Cholera toxin	Hypersecretion of electrolytes and water
	Toxin co-regulated pilus	Surface binding site receptor for bacteriophage CTXΦ; mediates bacterial adherence to intestinal mucosal cells
	Chemotaxis protein	Adhesin factor
	Accessory cholera enterotoxin	Increases intestinal fluid secretion
	Zonula occludens toxin	Increases intestinal permeability
	Neuraminidase	Modifies cell surface to increase GM <sub>1</sub> binding sites for cholera toxin




# *Vibrio cholera*

- Virulence of *V. cholerae* involved acquisition of first a sequence of genes including the **toxin co-regulated pilus (TCP)** on what is termed the **vibrio pathogenicity island (VPI-1)**, followed by infection with the **bacteriophage CTXΦ** that encodes the genes for the two subunits of **cholera toxin (ctxA and ctxB)**.
- TCP serves as the cell surface receptor for the bacteriophage, permitting it to move into the bacterial cell, where it becomes integrated into the *V. cholerae* genome.



# *Vibrio cholera*

- The cholera toxin is a **complex A-B toxin**. The active portion of the A subunit is internalized and interacts with G proteins that control adenylate cyclase, leading to the catabolic conversion of adenosine triphosphate (ATP) to cyclic adenosine monophosphate (cAMP). This results in a hypersecretion of water and electrolytes.
- The resulting severe fluid and electrolyte loss can lead to dehydration, painful muscle cramps, metabolic acidosis (bicarbonate loss), and hypokalemia and hypovolemic shock (potassium loss), with cardiac arrhythmia and renal failure.
- The mortality rate is as high as 70% in untreated patients but less than 1% in patients who are promptly treated with replacement of lost fluids and electrolytes



# *Brucella*


- Gram-negative bacteria, named after David Bruce (1855–1931). They are small (0.5 to 0.7 by 0.6 to 1.5  $\mu\text{m}$ ), non-encapsulated, non-motile, facultatively intracellular coccobacilli.
- Brucellosis in people is mostly caused by four species of *Brucella* bacteria. Most people get it through close contact with infected animals, often cows or through consumption of contaminated animal products.
- The gastrointestinal tract is affected in about 70% of cases, including anorexia, abdominal pain, vomiting, diarrhea, constipation, hepatomegaly, and splenomegaly.
- Due to the long incubation period for *Brucella*, cases often don't have symptoms until weeks or months after exposure

# *Brucella*

- **Brucellosis** in people cannot be diagnosed by clinical symptoms alone as initial symptoms are non-specific and can vary between patients. Laboratory testing is required to confirm a diagnosis.
- Most commercial clinical laboratories run serological tests that can detect antibodies to Brucella.
- Once brucellosis has been confirmed by lab testing, start treatment immediately to help prevent chronic infection-associated consequences

# Anaerobic Gram-Negative Rods

- These anaerobes are the **predominant bacteria on most mucosal surfaces, outnumbering aerobic bacteria 10- to 1000-fold.** Despite the abundance and diversity of these bacteria, most infections are caused by relatively few species
- ***Bacteroides*, *Fusobacterium*, *Parabacteroides*, *Porphyromonas*, and *Prevotella***
- Characteristically, *Bacteroides* growth is **stimulated by bile**. Other anaerobic gram-negative rods are fastidious. *Bacteroides* species are **pleomorphic in size and shape** and resemble a mixed population of organisms in a casually examined Gram stain.
- *Bacteroides* have a typical gram-negative cell wall structure, which can be surrounded by a **polysaccharide capsule**
- ***Bacteroides* LPS has little endotoxin activity**, probably due to **structural differences to pathogen LPS**.



**FIGURE 31-13** Growth of *Bacteroides fragilis* on *Bacteroides* bile-esculin agar. Most aerobic and anaerobic bacteria are inhibited by bile and gentamicin in this medium, whereas the *B. fragilis* group of organisms is stimulated by bile, resistant to gentamicin, and able to hydrolyze esculin, producing a black precipitate.

# Anaerobic Gram-Negative Rods

## Stomach $10^2$


*Lactobacillus*  
*Candida*  
*Streptococcus*  
*Helicobacter pylori*  
*Peptostreptococcus*

## Duodenum $10^2$

*Streptococcus*  
*Lactobacillus*

## Jejunum $10^2$

*Streptococcus*  
*Lactobacillus*



## Proximal ileum $10^2$

*Streptococcus*  
*Lactobacillus*

## Distal ileum $10^8$

*Clostridium*  
*Streptococcus*  
*Bacteroides*  
*Actinomycetaceae*  
*Corynebacteria*


## Colon $10^{12}$

*Bacteroides*  
*Clostridium*  
*Bifidobacterium*  
*Enterobacteriaceae*


# Anaerobic Gram-Negative Rods

- To cause disease, *Bacteroides fragilis* in the resident flora are able to **spread by trauma or disease from the normally colonized mucosal surfaces to sterile tissues or fluids**
- Infections are **usually polymicrobial**.
- **Respiratory Tract Infections**, Nearly half of the **chronic infections of the sinuses and ears**, and virtually all periodontal infections, involve **mixtures of gram-negative anaerobes**, with *Prevotella*, *Porphyromonas*, *Fusobacterium*, and **non-fragilis Bacteroides** the most commonly isolated.
- Strains of enterotoxigenic *B. fragilis* that cause diarrheal disease produce a **heat-labile zinc metalloprotease toxin (B. fragilis toxin)**. This toxin causes morphologic changes of the intestinal epithelium via F-actin rearrangement, with the resultant stimulation of chloride secretion and fluid loss. *B. fragilis* can produce a **self-limited watery diarrhea**.
- **Bacteremia**, Anaerobes were at one time responsible for more than 20% of all clinically significant cases of bacteremia; however, these organisms now cause 3% to 10% of such infections.

# Anaerobic Gram-Negative Rods



**FIGURE 31-11** Liver abscesses caused by *Bacteroides fragilis*.





**FIGURE 31-12** Synergistic polymicrobial infection involving *Bacteroides fragilis* and other anaerobes. The infection started at the scrotum and rapidly spread up the trunk and down the thighs, with extensive myonecrosis.

- **Intraabdominal Infections**, Anaerobes are recovered in **virtually all** of these infections, with ***B. fragilis* the most common organism**.
- **Skin and Soft-Tissue Infections**, *B. fragilis* is the organism most commonly associated with significant disease, usually in immunocompromised patients.

# Mycoplasma

- *Mycoplasma* and *Ureaplasma* organisms are the **smallest free-living bacteria**. They are unique among bacteria because they **do not have a cell wall** and their cell membrane contains **sterols**.
- The mycoplasmas form **pleomorphic shapes** varying from 0.2 to 0.3  $\mu\text{m}$  coccoid forms to rods 0.1 to 0.2  $\mu\text{m}$  in width and 1 to 2  $\mu\text{m}$  long.
- *M. pneumoniae* is a **strict human pathogen**.



# Mycoplasma

- **Respiratory disease** (e.g., tracheobronchitis, pneumonia) caused by *M. pneumoniae* occurs worldwide throughout the year.
- Exposure to *M. pneumoniae* typically results in **asymptomatic carriage**. The most common clinical presentation of *M. pneumoniae* infection is **tracheobronchitis**.
- Pneumonia (referred to as primary **atypical pneumonia**) can also develop, with a patchy bronchopneumonia seen on chest radiographs.
- *M. genitalium* can cause **nongonococcal urethritis (NGU)** and **pelvic inflammatory disease**.

Organism	Site	Human Disease
<i>Mycoplasma pneumoniae</i>	Respiratory tract	Tracheobronchitis, pharyngitis, pneumonia, secondary complications (neurologic, pericarditis, hemolytic anemia, arthritis, mucocutaneous lesions)
<i>Mycoplasma genitalium</i>	Genitourinary tract	Nongonococcal urethritis, pelvic inflammatory disease

# Mycoplasma

- **Microscopy** is of **no diagnostic value** because mycoplasmas stain poorly with the Gram stain. Likewise, antigen tests have poor sensitivity and specificity and are not recommended.
- The most sensitive diagnostic tests are **PCR amplification** tests of species-specific gene targets.
- Absence of the cell wall renders the mycoplasmas resistant to penicillins, cephalosporins, vancomycin, and other antibiotics that interfere with synthesis of the cell wall.

## Further reading:

- Murray - Medical Microbiology 8th Edition  
Section 4: Bacteriology  
Chapter 26:  
Chapter 28:  
Chapter 33: