

L (1) - The digestive system in the head and neck.

**Written by Sara Alkhateeb (23)
Reviewed by Mayar Khader**

1- The Mouth (Oral Cavity) - Anterior opening

The mouth (oral cavity) has **two** openings:


- ★ - **Anterior opening:** Between the upper and lower lips, bounded by the orbicularis oris muscle.
- **Posterior opening:** The oropharyngeal isthmus or fauces, which leads into the pharynx (oropharynx).

1- Lips

- The lips are two fleshy folds surrounding the oral orifice.
- **Externally** covered by skin, **internally** lined with mucous membrane.
- Essential for speech articulation, especially bilabial sounds like "p" and "m".
- Additional muscles radiate from the lips into the face.
- Contains labial blood vessels, nerves, connective tissue, and many small salivary glands.


2- Orbicularis oris muscle

- A circular striated muscle that acts as a sphincter, allowing actions like closing the mouth & whistling.
- Supplied by the facial nerve. Thus, ☹ **Injury to the facial nerve** results in:
 - 1 Inability to whistle (due to paralysis of orbicularis oris).
 - 2 Flattened buccinator muscle, making it hard to blow (not functional anymore).
 - 3- Drooling of saliva at the angle of the mouth during chewing due to lack of lip control.



Bounding Boundaries: The Oral Architecture

The mouth is a bounded physical space with distinct anterior and posterior gates.



1- The Mouth (Oral Cavity)

➤ Relate to HISTOLOGY

The Lip Consists of Three Main Zones:

1 Cutaneous (Outer) Zone




- Composed of keratinized stratified squamous epithelium (true skin).
- Contains hair follicles, sebaceous glands, and sweat glands.
- Hair is more prominent in males.

2- Mucosal (Inner) Zone

- Lined with non-keratinized stratified squamous epithelium.
- Contains labial glands (minor salivary gland), embedded within the mucosa and covered by the mucosal lining.



3- Red Zone (Vermilion or Transitional Zone)

- A type of modified skin (stratified squamous epithelium).
- Modified? since it Lacks hair follicles, sebaceous glands, and sweat glands.
- Appears red due to rich vascularity (blood vessels close to the surface).
- Highly sensitive due to the abundant nerve terminals.



1- The Mouth (Oral Cavity)

- **The philtrum:** is a shallow vertical groove in the midline of the upper lip. Formed by the fusion of the right and left medial nasal prominences on the outer surface of the upper lip.
- **Superior and Inferior Labial frenulae:** are mucosal folds that connect the inner surface of the upper/lower lips to the gums.



2- The Mouth Cavity – posterior opening


The mouth (oral cavity) extends from the lips to the pharynx. = two openings.

★ The **Posterior opening**: The oropharyngeal isthmus or fauces, which leads into the pharynx



Boundaries of the Oropharyngeal Isthmus:

- **Roof:** Formed by the soft palate and the uvula.
- **Floor:** Formed by the posterior third of the tongue, which contains lymphoid tissue known as the lingual tonsil (named due to its shape).
- **Lateral Walls:** Contain the palatine tonsils, which are commonly inflamed in children 😞 (**tonsillitis**) due to their role in filtering bacteria, viruses, and foreign bodies. These tonsils lie in a tonsillar fossa between two mucosal folds:

Anterior fold: The palatoglossal fold, containing the palatoglossus muscle.



Posterior fold: The palatopharyngeal fold, containing the palatopharyngeus muscle



3- Mouth divisions

The mouth is divided into the vestibule and the mouth cavity proper.

Vestibule > Space outside the closed teeth




- Slit-like space **located** between:

- **Anteriorly**: between the lips
- **Laterally/ externally** : bordered by the cheeks
- **Medially/ internally** : bordered by the gums and teeth

- **Function:**

1- The parotid duct opens on a small papilla in the vestibule opposite the upper second molar tooth, Saliva secreted here enters the oral vestibule, then passes into the mouth proper behind the last molar.

2- It communicates with the exterior through the oral fissure (the gap between the lips). When the jaws are closed, it still **communicates with the mouth proper behind the third molar tooth on each side.**





3- Mouth divisions

mouth cavity proper > cavity inside closed teeth

The mouth proper has a roof and a floor.

- The **roof** of the mouth is formed by the hard palate in front and the soft palate behind.

- The **floor** is formed largely by the anterior two thirds of the tongue



- The **lateral side** is formed by closed teeth (gums are the tissues that surround the teeth and help hold them in place, essentially acting as a socket for the teeth).

□ There are two **types of mucous membranes** based on their underlying connective tissue:

1 **Soft and Elastic (CT) Mucosa:** Found in: Mucosa of lips Floor of the mouth & Underside of the tongue.

2 **Dense (CT) Mucosa:** Found in: Hard palate, gums (gingiva), surrounding the teeth

□ Mouth **blood supply:** branch of facial artery & branch of lingual Artery.



4- Sensory Innervation of the Mouth

Roof of the Mouth (Hard Palate)

- Greater palatine nerve
- Nasopalatine nerve


(Both are branches of the maxillary V2 division of the trigeminal nerve -5th CN)

Floor of the Mouth:

General Sensation (touch, temperature, pain)

- **Lingual nerve**

(Branch of the mandibular V3 division of the trigeminal nerve - CN)



□ Taste areas

- Posteriorly (Circumvallate): bitter
- Laterally: sour and salty
- Anteriorly : sweet

Special Sensation (Taste)

- **Chorda tympani nerve**

(Branch of the facial nerve) These fibers carry taste sensations from taste buds located on the dorsum of the tongue

Cheek

Sensory (Outer Surface (Skin) and Inner Mucosa)

- **Buccal nerve**


(Sensory branch of the mandibular division of the trigeminal nerve - CN V3)

Motor (Buccinator muscle)

- **Buccal branch of the facial nerve**

(Extra slide) Recall :

* NERVE SUPPLY *





TeachMeAnatomy

- Ophthalmic (V1)
- Maxillary (V2)
- Mandibular (V3)

5- The teeth

- The gingivae (gum) are specialized regions of the oral mucosa that surround the teeth.
- They are surrounded by dense connective tissue.




6- The teeth

-	Deciduous Teeth (Milk Teeth)	Permanent Teeth
# number	20 teeth ; 10 in each jaw.	32 teeth ; 16 in each jaw.
Types per jaw	(4) Incisors, (2) Canines & (4) Molars	(4) Incisors, (2) Canines, (4) premolars & (6) Molars
Eruption timeline	Begins around 6 months of age, all 20 teeth typically erupt by 2 years of age.	Begin to erupt at the age of 6 and continue until around 12 years. The third molar, Aka *wisdom tooth, is the last to erupt and typically appears between the ages of 17 and 30.
Eruption Order	Lower jaw teeth usually erupt before the upper jaw teeth.	
Notes	-	*They're called "wisdom teeth" because they appear later in life. Their eruption can be associated with various issues. In many cases, they may remain unerupted (impacted) and can sometimes lead to infection.

7- The tongue

- ❑ The tongue is a muscular organ made of striated “muscle” covered by “mucous membrane”.
- ❑ A **median fibrous septum** divides the tongue into symmetrical right and left halves, each with paired muscles. – *This division is significant as it allows for the coordinated action of the intrinsic and extrinsic muscles (muscular imp.)*
- ❑ The tongue is also anatomically divided into an anterior two-thirds and a posterior one-third by the **sulcus terminalis and the foramen cecum**. – *division is significant as it corresponds to different types of mucous membrane lining the tongue (mucous membrane importance)*



8- Mucous membrane of the tongue

□ The tongue is a muscular organ made of striated muscle covered by “**mucous membrane**”.

➤ **Relate to HISTOLOGY**

- The mucous membrane of the tongue is divided into two parts: the dorsal (upper) surface and the ventral (lower) surface.

★ • **The upper (dorsal) surface of the tongue:**

- lined with stratified squamous **para-keratinized** epithelium.
- It was originally keratinized, but due to digestion and secretion, it becomes **injured** and changes to para- keratinized.

• **The lower (ventral) surface**

- Covered by stratified squamous **non-keratinized** epithelium.

The mucous membrane of the upper surface of the tongue can be divided into anterior and posterior parts, by a V-shaped sulcus called “**sulcus terminalis**”. Where the apex of the sulcus projects backward and is marked by a small pit, called “**foramen cecum**”.

8- Mucous membrane of the tongue

1- upper surface of the tongue

Anterior 2/3


Filled with **taste buds**, AKA lingual papillae, including filiform, fungiform, and foliate & vallate papillae. Where filiform are the most abundant.

posterior 1/3

Devoid of papillae but has an irregular surface caused by the presence of underlying **lymphoid tissue** (lingual tonsil / follicle).

2- inferior surface of the tongue

- The mucous membrane on the inferior surface of the tongue is reflected from the tongue to the floor of the mouth.
- In the **midline** anteriorly, the undersurface of the tongue is connected to the floor of the mouth by a fold of mucous membrane: **frenulum of the tongue**.
- On the **lateral** side of the frenulum, the **deep lingual vein** can be seen through the mucous membrane. The lingual artery and lingual nerve lie deeper and more medially, and are not typically visible, but they run in close relation to the vein beneath the mucosa.



8- Mucous membrane of the tongue

❑ Taste buds


Taste buds are specialized sensory structures found mostly on the dorsal (upper) surface of the tongue. They are responsible for detecting different taste sensations like sweet, salty, sour, bitter, and umami (savory).

- **Fungiform Papillae** Scattered across the anterior part of the tongue.
- **Foliate Papillae** Lateral edges of the posterior part of the anterior two-thirds.
- **Circumvallate (Vallate) Papillae** Arranged in a V-shape just anterior to the sulcus terminalis. Note: exception! they sit anteriorly but are innervated posteriorly due to their embryological origin. - Refer to slide (25).

✗ **Filiform Papillae** Most numerous. Do NOT contain taste buds. Function more for tactile sensation (touch/texture).

❑ Taste areas

- Posteriorly (Circumvallate): bitter
- Laterally: sour and salty
- Anteriorly : sweet



9- Muscles of the tongue

- The tongue is a muscular organ made of striated “**muscle**” covered by mucous membrane.

It has both

-	intrinsic muscles	extrinsic muscles
Consists of:	longitudinal, transverse, & oblique fibers	styloglossus, genioglossus , palatoglossus, & hyoglossus muscles)
Attachment	They are confined to the tongue and are not attached to bone	muscles are attached to bones and the soft palate
Action	control & alter tongue shape	control its movement and connect it to surrounding structures like the styloid process, soft palate, mandible, and hyoid bone.
Nerve supply	Hypoglossal nerve except palatoglossus supplied by cranial accessory of vagus nerve.	



10- Extrinsic muscles of the tongue

-	Origin	Insertion	Action	Nerve supply
styloglossus	Styloid process of temporal bone	base of the tongue posteriorly	Draws tongue upward and backward	Hypoglossal nerve (Cranial Nerve XII) Except palatoglossus supplied by cranial part of the accessory nerve (CN XI) via the vagus nerve (CN X).
genioglossus	Superior genial (tubercle) of the spine of mandible	base of the tongue posteriorly	the ONLY muscle that protrudes the tongue out of the mouth.	
hyoglossus	hyoid bone	base of the tongue	Depresses tongue	
palatoglossus	Palatine aponeurosis	Side of tongue	Pulls roots of tongue upward and backward, narrows oropharyngeal isthmus	

Notice how the muscles attach the tongue to the styloid process and the soft palate above and to the mandible and the hyoid bone below.

- Slide 22 -


10- Extrinsic Muscles of The tongue



11- Movements of the Tongue

❑ **Protrusion** (Sticking the tongue out)

Muscle involved: **Genioglossus** (both sides working together)



Normally: tongue remains straight when protruded outside the mouth because both genioglossus muscles (on the right and left sides) contract symmetrically. Thus ☹ **Injury to the hypoglossal nerve** results in **ipsilateral** tongue paralysis and deviation toward the affected side.

❑ **Retraction** (Pulling the tongue back)

Muscles involved: Styloglossus & Hyoglossus (both sides acting together)

❑ **Depression** (Lowering the tongue)

Muscle involved: Hyoglossus (both sides working)

❑ **Retraction and Elevation of the Posterior Third**

Muscles involved: Styloglossus & Palatoglossus (both sides acting together).


Function: Pulls the back of the tongue up and back, important during swallowing

(Extra slide)

NOT REQUIRED:

Injury to the hypoglossal nerve


Normally:




The genioglossus muscle on each side pulls the tongue forward and slightly toward the opposite side. **The right genioglossus** pulls the tongue forward and to the left. The **left genioglossus** pulls the tongue forward and to the right. So, when both genioglossus muscles contract with equal strength Their opposite directional forces cancel each other out. As a result, tongue is protruded straight forward, staying in the midline.

VS

Injury:



Our case is lower motor neuron lesions ★ (injury to the hypoglossal nerve), the tongue deviates the same side as the lesion.




right genioglossus muscle is paralyzed. The **left genioglossus** works normally and pushes the tongue forward and slightly to the right. → Deviation occurs due to unopposed action.

12- Sensory innervation of the tongue


➤ Relate to EMBRYOLOGY

During embryological development, the anterior two-thirds of the tongue arises from the first pharyngeal arch, while the posterior third comes from the third pharyngeal arch.



As a result, their nerve supply differs:


- The **posterior third** is innervated by the **glossopharyngeal nerve** (CN IX) for both general sensation and taste.
- The **anterior two-thirds** of the tongue receives:
 - General sensation from the **lingual nerve**.
 - Taste sensation from the **chorda tympani**.



12- Sensory innervation of the tongue

➤ Relate to EMBRYOLOGY

- Recall: The circumvallate papillae, which are responsible for sensing bitter taste, are located near the sulcus terminalis.– slide 18
- Embryologically, these papillae **belong to the posterior one-third** of the tongue, even though **anatomically** they appear in the anterior part.
- Because of their embryological origin, they are innervated by the **glossopharyngeal nerve**. This differs from the rest of the anterior 2/3 of the tongue, which is innervated by the chorda tympani.
- The posterior one-third of the tongue is largely made of lymphoid tissue (lingual tonsil), and normally lacks taste buds, but the circumvallate papillae still provide taste sensation in this region.



13- Blood supply & Lymphatics of the tongue

■ Blood Supply of the Tongue

The tongue receives arterial blood from branches of the external carotid artery, including:


- **Lingual artery** (main supply)
- **Tonsillar branch of the facial artery**
- **Ascending pharyngeal artery**

■ Venous Drainage

The veins of the tongue drain into the **internal jugular vein**.

■ Lymphatic Drainage

- Tip of the tongue drains into the **submental lymph nodes**.
- “Others” drains into **submandibular**.
- All lymph from these nodes eventually drains into the **deep cervical lymph nodes**.



* OTHERS? → should be Lateral anterior two-thirds of the tongue.


14- Salivary glands – introduction

Types

Major Salivary Glands

- **Parotid gland:** Has serous secretion and two capsules.
- **Submandibular gland:** Has mixed secretion (serous + mucous) and one capsule.
- **Sublingual gland:** Has mostly mucous secretion and one capsule.

Minor Salivary Glands

- 
- **Labial** glands (within the lips)
 - **Lingual** glands (in the tongue)
 - **Palatal** glands (in the palate)

Numerous and mostly mucous glands.

○ Innervation of Salivary Glands

Sensory (for general sensation)

Sympathetic (for *vasoconstriction of blood vessels)

Parasympathetic (for *secretomotor function)


*Parasympathetic: Directly stimulates saliva secretion.

*Sympathetic: Does not directly inhibit secretion but indirectly reduces it by **vasoconstricting** the blood vessels → leading to less blood supply, which affects secretion.

16- Parotid gland

- Largest salivary gland, Mostly serous acini.

Consists of 1- **glandular part** & 2- **ductal system**.



The parotid duct emerges from the anterior border of the gland → runs anteriorly over the external surface of the masseter muscle → turns medially, pierces the buccinator muscle, → opens into the oral cavity adjacent to the crown of the second upper molar tooth.

- It is approximately 5 cm long.
- Surface anatomy: one fingerbreadth below the zygomatic arch.


- The parotid gland is located anterior to the ear, and it overlies both the masseter muscle and the sternocleidomastoid muscle.
- The base of the gland lies just beneath the skin, while the apex points toward the pharynx.

16- Parotid gland

- The parotid gland lies in the “parotid bed” (anatomical space upon which the gland rests).

- which is anatomically **bounded** by:

1. Posteriorly: **sternocleidomastoid** muscle.
2. Anteriorly : **ramus of the mandible**.
3. Superiorly : **external acoustic meatus & posterior part of the zygomatic arch**.




- Additionally, the parotid bed includes several other **structures** that the gland rests upon:

1. The **styloid process of the temporal bone**.
2. Muscles attached to the styloid process, such as the **stylohyoid** muscle.
3. The **posterior belly of the digastric muscle**.
4. The **deep cervical fascia**.
5. The **last four cranial nerves (CN IX–XII)**.

Common question:


16- Parotid gland



Main structure	Others	<p>★ parotid gland overlies major blood vessels:</p> <ul style="list-style-type: none"> - Internal jugular vein. - Common carotid artery which bifurcates into the external (does supply the gland) & internal carotid arteries (doesn't supply the gland)
<p>1 Facial nerve (CN VII) “extra cranial part” (most superficial)</p> <p>2 Retromandibular Vein (intermediate)</p> <p>3 external carotid artery (deepest)</p>	<p>4 Parotoid lymph nodes</p> <p>5 Auriculotemporal Nerve</p> <p>6 Posterior auricular artery</p>	

16- Facial Nerve in the parotoid gland

- The most superficial structure within the parotid gland is the facial nerve & its five terminal branches, they pass through the gland but do not supply it functionally!
- The **main trunk** of the facial nerve enters the parotid gland and divides it into **superficial & deep lobes**.
- Whereas the **branches** arise within the gland and fan out to **innervate the muscles of facial expression**:
 1. **Temporal** branch → innervates the orbicularis oculi
 2. **Zygomatic** branch
 3. **Buccal** branch
 4. Marginal **mandibular** branch → innervates muscles near the mandible
 5. **Cervical** branch → innervates the platysma




During ☹️ **parotid gland surgery (e.g. tumor removal), the facial nerve is at risk of injury**. This is a primary surgical concern. Thus, On the first day after surgery, the doctor typically assesses facial nerve function by asking the patient to perform actions involving facial muscles:

- Close the eyes → tests the temporal branch (orbicularis oculi)
- Blow out the cheeks
- Check for drooling of saliva


Recall slide 5

16- Structures with the parotid gland


- The **retromandibular vein**, lies deep to the facial nerve, formed by the union of the superficial temporal and maxillary veins within the parotid gland.
- The **external carotid artery**, along with its terminal branches (superficial temporal and maxillary arteries), lies deeper within the gland and provides the arterial supply to the region.
- The parotid gland contains intraglandular lymph nodes, commonly referred to as the **parotid lymph nodes**.




Anatomy of the Salivary Glands



16- Parotoid gland




- Recall: The parotid gland has 2 capsules.
- The parotid gland is divided into **lobes** and **lobules**.
- The lobes are larger sections of the gland, while the lobules are smaller subunits within the lobes. They are separated by **connective tissue septa**, which extend inward from the **inner capsule** of the gland.
- Within each lobule, there are secretory acini, which are small glandular structures that produce saliva. These acini drain into small interlobular ducts, which are smaller tubes that collect the secretions from multiple acini. These interlobular ducts then merge to form the **main parotid duct**, which is the primary duct responsible for transporting saliva from the gland to the oral cavity.
- The **outer capsule** of the parotid gland is made of parotid fascia, a type of connective tissue that surrounds the gland. This fascia is derived from the deep cervical fascia.
- ☹️ **Mumps (أبو دغيم)** is a **viral infection** that primarily affects the parotid gland. The virus causes inflammation and swelling of the gland, which can lead to discomfort and pain. The fibrous capsule of the parotid gland, which is tough and not very elastic, restricts the gland's ability to expand. This limitation results in severe pain due to the pressure buildup inside the gland as it swells.



16- Parotoid gland Innervation

The **parasympathetic innervation** of the parotid gland originates from: **inferior salivatory nucleus** in the medulla oblongata. The **glossopharyngeal nerve** (cranial nerve IX) emerges from this nucleus and gives rise to the **tympanic nerve**, which branches into the **lesser petrosal nerve** at the tympanic membrane. The lesser petrosal nerve carries preganglionic parasympathetic fibers to the **otic ganglion**, located just below the foramen ovale. The postganglionic parasympathetic fibers then travel via the **auriculotemporal nerve** (a branch of **Mandibular**) to reach and innervate the parotid gland and provides **sensory** innervation to the gland.




Note: **Postganglionic sympathetic fibers** originates from the superior cervical ganglion in the neck and reach the gland via a plexus around the external carotid artery.

Parasympathetic secretomotor supply arises from

17- Submandibular gland

- Recall: A mixed salivary gland, composed of both serous and mucous acini.
Consists of 1- **glandular part** & 2- **ductal system**.




(Wharton's Duct)

- Arises from the anterior end of the deep part of the gland.
 - It travels forward in the floor of the mouth and opens at the base of the tongue, on a small papilla at the side of the lingual frenulum (the submandibular papilla).
- It is located beneath the lower border of the body of the mandible, occupying the submandibular (digastric) triangle, which is bounded by the anterior and posterior bellies of the digastric muscle.
 - The gland is divided into two parts **by the mylohyoid muscle**:
 - **Superficial** part: Lies superficial (inferior) to the mylohyoid.
 - **Deep** part: Lies deep (superior) to the mylohyoid, beneath the mucous membrane of the floor of the mouth and adjacent to the side of the tongue.
 - * The two parts are continuous around the posterior border of the mylohyoid muscle.

18- Sublingual gland


- It is the **smallest** major salivary gland, enclosed in a thin mucosal capsule.
- It contains both serous and mucous acini, but **mucous acini predominate**.
- The sublingual gland lies beneath the tongue, embedded in the floor of the mouth, specifically under the mucous membrane of the sublingual fold, and is located close to the lingual frenulum.
- It has **8–20 small ducts** that open into the mouth on the summit of the sublingual fold, and in some cases, some ducts join the submandibular duct.
- Medially related to the ***lingual nerve, submandibular duct, and genioglossus muscle**.



* Because the secretion is viscous (thick mucus), the gland needs multiple small ducts (*unlike the parotid or submandibular glands*) to drain efficiently without clogging or pooling. This way, the secretion is evenly spread into the floor of the mouth — especially helpful during speaking, chewing, and swallowing!

Submandibular & sublingual glands


* Origin of Mylohyoid muscle separates the submandibular gland from the sublingual gland



19- Submandibular & Sublingual gland Innervation

1. Parasympathetic & Sensory Innervation

Pre-Ganglionic Phase	<ul style="list-style-type: none"> • Originates from the superior salivatory nucleus in the medulla oblongata, carried by the facial nerve. • The facial nerve gives rise to the chorda tympani nerve, which carries preganglionic parasympathetic fibers as well as taste fibers from the anterior two-thirds of the tongue. • The chorda tympani joins the lingual nerve in the infratemporal fossa.
Synapse	<ul style="list-style-type: none"> • At: submandibular ganglion, which is located within the submandibular triangle, between the mylohyoid and the hyoglossus muscles. * Note: The chorda tympani fibers travel with the lingual nerve to reach the ganglion; they do not run independently, But their fibers remain distinct.
post-Gangl.	<ul style="list-style-type: none"> • Postganglionic parasympathetic fibers either directly innervate the submandibular and sublingual glands or re-enter the lingual nerve to reach their target glands.




* The lingual nerve provides general sensory fibers. It also carries parasympathetic fibers & taste fibers from the chorda tympani to the submandibular gland.

- **Preganglionic parasympathetic: chorda tympani**
- **Sensory and taste fibers: lingual nerve**

19- Submandibular & Sublingual gland Innervation


1. Sensory Innervation – comp.

- The **lingual nerve** has a **triple relation** to the **submandibular duct** it passes **lateral**, then **inferior**, and finally **medial** to it.





2. Sympathetic Innervation

- Postganglionic sympathetic fibers: arise from the **superior cervical ganglion** and reach the glands via a plexus around the lingual artery.



20- Salivary glands – To sum up:



Mylohyoid muscle.

- separates the superficial part of the submandibular gland from its deep part.
- It also separates the submandibular gland from the sublingual gland.