



# Cranial Cavity Part 2

---

**Dr. Ahmed Salman**

**Associate professor of anatomy & embryology**

# **Dural venous sinuses**

---



## Dural Venous Sinuses

- They are wide venous channels which lie between the outer periosteal and meningeal layers of dura **EXCEPT** the **inferior sagittal and straight sinuses** which lie within dural folds
- They are lined by endothelium
- No muscle tissue in their wall .
- They have no valves
- They communicate with the veins outside the skull by the *valueless* emissary veins.
- They drain bones of the skull, meninges, C.S.F., end brain into the I.J.V.

# Classification

Unpaired group	Paired group
<ol style="list-style-type: none"><li>1. Superior sagittal sinus</li><li>2. Inferior sagittal sinus</li><li>3. Straight sinus</li><li>4. Occipital</li><li>5. Basilar plexus</li><li>6. Intercavernous</li></ol>	<ol style="list-style-type: none"><li>1. Sphenoparietal</li><li>2. Cavernous</li><li>3. Superior petrosal</li><li>4. Inferior petrosal</li><li>5. Transverse</li><li>6. Sigmoid.</li></ol>

## Superior Sagittal Sinus :


It runs in the superior border of the falx cerebri and near the internal occipital protuberance, it deviates (usually) to the **right** to become the **right transverse** sinus.


It receives :

**1- Superior cerebral veins** which are thin-walled

Rupture of these veins cause subdural hemorrhage with compression of the brain.

**2- Arachnoid granulations** which filter C.S.F. into the venous circulation





## Inferior Sagittal Sinus :

- ❖ It lies in the posterior 2/3 of the lower (free) border of falx cerebri.
- ❖ It joins the great cerebral vein to form the straight sinus.


## Straight Sinus :

- ❖ It lies at the junction between falx cerebri and the tentorium cerebelli.
- ❖ It deviates usually to the **left** to become the **left** transverse sinus.


**Inferior Sagittal Sinus**

**JOIN**

**Great cerebral vein**




**Straight**




**Left transverse sinus**

**Superior Sagittal Sinus**



**Right transverse sinus**






## Inferior Sagittal Sinus

© www.kenhub.com






## **Straight Sinus**

## The Cavernous Sinuses :

It lies between the 2 layers of the dura on the side of the body of the sphenoid bone

It extends from superior orbital fissure anteriorly to apex of petrous bone posteriorly





## **The sinus receives :**

- 1-The superior ophthalmic vein
- 2-The inferior ophthalmic vein
- 3-The central vein of the retina
- 4-Superficial middle cerebral vein
- 5- Sphenoparietal sinus

## **The sinus drains posteriorly into:**

1. The transverse sinus through the superior petrosal sinus
2. Internal jugular vein through Inferior petrosal sinus






## Structures traverse the cavernous sinus :


- The inferomedial wall of the sinus is traversed by the 3rd part of the internal carotid artery and the Abducent N. (6<sup>th</sup> cranial nerve) , inferolateral to the artery .
- The lateral wall is traversed (from above downwards) by the oculomotor, trochlear nerves , ophthalmic and maxillary nerves.

All the above structures are *separated* from the blood within the sinus by the sinus endothelium.



# Cavernous Sinus

## Coronal Section



## **The Transverse Sinuses :**

- Each runs in the transverse sulcus between the 2 layers of the attached margin of tentorium cerebelli
- The right sinus is continuous with the superior sagittal sinus; the left with the straight sinus.
- Each terminates by becoming the sigmoid sinus.

## **The Sigmoid Sinuses :**

- This S-shaped sinus is the direct continuation of the transverse sinus.
- It begins behind the base of the petrous temporal bone and passes through the posterior compartment of the jugular foramen where it continues as the Internal jugular vein.


## The confluence of sinuses Or torcula :

**Site :** internal occipital protuberance


It is the site of communications between 5 venous sinuses

1. Superior sagittal
2. Straight
3. Occipital
4. Right and left transverse sinuses.

Thrombosis in one sinus will spread to the other sinuses in the confluence.



## Cerebral veins - Anterior view



### Legend

1. Internal jugular vein
2. Sigmoid sinus
3. Transverse sinus
4. Superior sagittal sinus
5. Superior cerebral veins
6. Vein of Labbe
7. Torcula

## **Middle meningeal Artery**

- It is a branch from the first part of the maxillary A. in the infratemporal fossa.
- It enters the skull through the foramen spinosum.
- It runs intracranial anterolateral between the 2 layers of the dura
- 2 cm above the midpoint of the zygomatic arch, the artery divides into frontal and parietal branches.


1-The frontal branch : (is the larger) crosses the greater wing of the sphenoid and ascends towards the pterion.


Here, it is related medially to the motor area of the cerebral cortex

2-The parietal branch : runs backwards on the squamous temporal bone, and is related medially to the superior temporal gyrus of the cerebral hemisphere.

### Branches of middle meningeal

- ❖ To the inner table and diploe of the skull bones and dura mater.
- ❖ It gives the superior tympanic A. and petrosal branches to the tympanic cavity.







## Epidural Hemorrhage

- Results from injuries to the meningeal arteries or veins.
- The most common artery to be damaged is the **anterior division of the middle meningeal artery**
- Arterial blood is located between the skull and dura
- The intracranial pressure rises, and the enlarging blood clot exerts local pressure on the underlying motor area
- Lucid interval (no symptoms) for a few hours followed by death (“talk and die syndrome”)

# Epidural Hemorrhage



**Look for this website :**

[https://www.radiologymasterclass.co.uk/gallery/ct\\_brain/ct\\_brain\\_images/extradural\\_contraco\\_up\\_ct\\_brain](https://www.radiologymasterclass.co.uk/gallery/ct_brain/ct_brain_images/extradural_contraco_up_ct_brain)


## Treatment

- The burr hole through the skull wall should be placed 2.5 to 4 cm above the midpoint of the zygomatic arch to ligate or plug the torn artery or vein.



## Subdural hemorrhage


- ❑ Is caused by a violent shaking of the head (e.g. car accident)
- ❑ The blood is located between the dura and arachnoid
- ❑ Blood accumulates slowly (days to weeks after trauma);
- ❑ No blood in the CSF after lumbar puncture.





**Epidural Hematoma**  
(Does Not Cross Suture Line)

**Subdural Hematoma**  
(Crosses Suture Line)



**Look for this website :**

[https://www.radiologymasterclass.co.uk/gallery/ct\\_brain/ct\\_brain\\_images/subdural\\_small\\_ct\\_brain](https://www.radiologymasterclass.co.uk/gallery/ct_brain/ct_brain_images/subdural_small_ct_brain)

## Subarachnoid hemorrhage :

- Extravasation of blood in Subarachnoid space usually arterial.
- Th cause traumatic as head truma or non traumatic like rapture arterial aneurysm
- This hemorrhage resulted in meningeal irritation ,severe headache ,neck stiffness and loss of consciousness

### Look for this website :

[https://www.radiologymasterclass.co.uk/gallery/ct\\_brain/ct\\_brain\\_images/subarachnoid\\_haemorrhage\\_sah\\_ct\\_brain](https://www.radiologymasterclass.co.uk/gallery/ct_brain/ct_brain_images/subarachnoid_haemorrhage_sah_ct_brain)

## Cerebral hemorrhage

Is caused by rupture of the thin-walled a branch of **the middle cerebral artery**.

The hemorrhage involves the important centers


It produces hemiplegia on the opposite side of the body and may be affect speech

### Look for this website :

[https://www.radiologymasterclass.co.uk/gallery/ct\\_brain/ct\\_brain\\_images/intracerebral\\_haemorrhage\\_ich\\_ct\\_brain](https://www.radiologymasterclass.co.uk/gallery/ct_brain/ct_brain_images/intracerebral_haemorrhage_ich_ct_brain)

Epidural Hematoma

Subdural Hematoma



Intracerebral Hemorrhage

Subarachnoid Hemorrhage



Thank  
you

