



The background of the slide is a light pink color, populated with numerous stylized illustrations of blood cells. Red blood cells are depicted as red, biconcave discs of varying sizes. White blood cells are shown as larger, more complex shapes: some are purple with internal granules and a multi-lobed nucleus, while others are blue with a large, kidney-shaped nucleus and a scalloped outer edge. Small, yellow, bean-shaped cells are also scattered throughout the field.

Hematology Physiology

Fatima Daoud, MD, PhD.

The background of the slide is a light pink color, populated with numerous stylized illustrations of blood cells. Red blood cells are depicted as red, biconcave discs of varying sizes and orientations. White blood cells are shown as larger, spherical cells with purple or blue nuclei and granular cytoplasm. Platelets are represented as small, yellow, oval-shaped cells. The cells are scattered across the entire slide, creating a dense, microscopic-like pattern.

Reference books:

1. **Hall, John, E. and Michael E. Hall. Guyton and Hall Textbook of Medical Physiology (14th Edition).**
2. Lauralee Sherwood. Human Physiology: From Cells To Systems (9th Edition).
3. Gerard J. Tortora and Bryan Derrickson. Principles Of Human Anatomy & Physiology (15th Edition)

Hematology

Body Fluids






we divide body fluid compartments into :-

* ICF (Intracellular fluid)

* ECF (Extracellular fluid)

→ Plasma
→ interstitial fluid

There are differences in the electrolyte composition between these compartments.



(a) Distribution of body solids and fluids in average lean adult female and male



Hematology

Body Fluids

The main difference between intracellular compartments & Extracellular compartments is in:-

- ① Na^+
- ② K^+
- ③ Mg^{+2}
- ④ Cl^-
- ⑤ HPO_4^{2-}
- ⑥ SO_4^{2-}
- ⑦ Protein Anions.

Key:



- The average blood volume of adults is about **7–8%** of **body weight**, or **about 5 liters**.
- Blood is **connective tissue**.
- Blood is **denser** and **more viscous (thicker)** than water.
- The temperature of blood is **38°C**.
- **Slightly alkaline** pH ranging from **7.35** to **7.45** (average = **7.4**).
- The color **saturated (O_2)** → **bright red**
unsaturated (O_2) → **dark red**


Hematology

Blood and its components

Blood smear

Blood Smear

This an important approach for diagnosing Cancers, & Anemia (in the past, now we use more automated methods).




White blood cell
(leukocyte: neutrophil)

Blood plasma

Red blood cell
(erythrocyte)


Platelet (not an actual cell, it is a fragmentation of Megakaryocyte).

White blood cell
(leukocyte: monocyte)



Hematology

Blood and its components



are major difference between plasma & interstitial fluid

Hematology

Plasma Protein

1. Establishment of **colloid osmotic pressure**.

Drive the movement of plasma from the interstitial fluid into the plasma and prevents the leakage of fluid out of vascular system.

Key:


BHP = Blood hydrostatic pressure
IFHP = Interstitial fluid hydrostatic pressure
BCOP = Blood colloid osmotic pressure
IFOP = Interstitial fluid osmotic pressure
NFP = Net filtration pressure

Blood hydrostatic pressure pushes fluid out of capillaries (filtration)

Blood colloid osmotic pressure pulls fluid into capillaries (reabsorption)

* One of the main drivers of colloid oncotic pressure is albumin which is synthesised by the liver, so if there is any liver problem that inhibits the albumin synthesis, this is going to cause edema.

* Plasma proteins work as a transporters for lipid soluble substances.



	Net filtration at arterial end of capillaries (20 liters per day)	Net reabsorption at venous end of capillaries (17 liters per day)
Net filtration pressure (NFP)	$(BHP + IFOP) - (BCOP + IFHP)$	$(BHP + IFOP) - (BCOP + IFHP)$
	Pressures promoting filtration	Pressures promoting reabsorption
	Arterial end	Venous end
	$NFP = (35 + 1) - (26 + 0)$ $= 10 \text{ mmHg}$	$NFP = (16 + 1) - (26 + 0)$ $= -9 \text{ mmHg}$
Result	Net filtration	Net reabsorption

3 Liters.
absorbed back by lymphatic system to vascular circulation.

Hematology

Plasma Protein

2. Responsible for plasma's capacity buffer changes in pH.

Albumin

Synthesised from liver

Nonspecifically binds substances that are poorly soluble in plasma (bilirubin & fatty acids)

Fibrinogen

Is an inactive precursor for a clot's fibrin meshwork

Globulins

α & β

- Specifically bind poorly water-soluble substances (thyroid hormone, cholesterol, and iron).
- Involved in blood-clotting.
- Angiotensinogen.

γ

Antibodies/ immunoglobulins.
Body's defense mechanism.

[Immunity]

Transportation

Respiratory gases, nutrients, hormones, and metabolic wastes

Regulation

↑↑ buffer changes in pH
Acid-base balance, heat distribution, and fluid balance (through osmolarity).
↗ Vaso dilation.
↘ Vaso constriction.

Protection

Immune defense and prevention of blood loss

Hematology

Hemopoiesis (formation of blood cells)

Early fetal life

① Occurs in the **yolk sac** of an **embryo** and **later in** the **liver, spleen, thymus, and lymph nodes** of a fetus.
②

* Hemopoiesis' location throughout life:-


last gestational months- death

* **Red bone marrow** becomes the primary site of hemopoiesis in the, and continues as the source of blood cells after birth and throughout life.

Hematology

Hemopoiesis (formation of blood cells)

Red bone Marrow → located in the core of bones, but which bones?! it depends on age.



As we are getting older the Hemopoiesis process shifts from being happening mainly at the shaft of long bones towards their ends.

As we are hitting 20 y.o. the hemopoiesis process becomes happening mainly at flat bones.

it decreases with age.

Key:

- Progenitor cells
- Precursor cells or "blasts"
- Formed elements of circulating blood
- Tissue cells

Key:

- CFU-E Colony-forming unit—erythrocyte
- CFU-Meg Colony-forming unit—megakaryocyte
- CFU-GM Colony-forming unit—granulocyte macrophage

CFU are committed regenerating cells. it means colonies $\Rightarrow$ specialized

toward generating specialized cells

Pluripotent stem cell

little degree differentiation to

Myeloid stem cell

Lymphoid stem cell

Colony forming units

CFU-E

CFU-Meg

CFU-GM

Proerythroblast

Megakaryoblast

Eosinophilic myeloblast

Basophilic myeloblast

Myeloblast

Monoblast

T lymphoblast

B lymphoblast

Natural killer

NK lymphoblast

Nucleus ejected

Reticulocyte

Megakaryocyte

Red blood cell (erythrocyte)

Platelets

Eosinophil

Basophil

Neutrophil

Monocyte

T lymphocyte (T cell)

B lymphocyte (B cell)

Natural killer (NK) cell


Mast cell

Granular leukocytes

Macrophage

Agranular leukocytes

Plasma cell



- ✓ **Myeloid stem cells**
 - Give rise to red blood cells, platelets, monocytes, neutrophils, eosinophils and basophils
- ✓ **Lymphoid stem cells** give rise to
 - Lymphocytes and natural killer cells

- **Stem cells in bone marrow**
 - Reproduce themselves
 - Proliferate and differentiate

- **Formed elements do not divide once they leave red bone marrow**

*under physiological conditions
They have already been a specific type of WBCs which normally don't divide*

- Exception is lymphocytes


What causes stem cells to differentiate to a specific type of white blood cells?! inducers. (signals) / sometimes called differentiation inducer.

- **Growth inducers:** regulate growth and reproduction of the different stem. One of the growth inducers is **interleukin-3**, promotes growth and reproduction of virtually all the different types of committed progenitor cells. (**GM-CSF**, **G-CSF**, and **M-CSF**)
- **Differentiation inducers:** Each of these differentiation inducers causes one type of committed stem cell to differentiate one or more steps toward a final adult blood cell.
- Others: **Erythropoietin – RBCs**, **Thrombopoietin – platelets** (*Thrombocytes is the other name).*
- Signals from outside the marrow match production to the body's needs

Hematology

Packed Red Cell Volume

PCV is a laboratory test that is used to reflect the volume of RBC to the total blood volume
→ now it is automated.



Hematology

Hematocrit

- **Hematocrit/ Packed Red Cell Volume**


- Adult males: 40–54% (avg = 45%). *more RBCs Volume in men.*
- Adult females: 38–46% (avg = 42%) *memorise*

$$\text{Hematocrit} = \frac{\text{Height of RBCs}}{\text{Total height}} \times 100\%$$

→ Hematocrit is a ratio
no direct indication on ↑ number of RBCs.

* **Ratio!!**

why?! Testosterone activates Hemopoiesis + menstrual cycle in women.




Hematology


Hematocrit

↑ (RBC)

Polycythemia



Anemia



(↓ RBC)

- * An increased hematocrit can reflect more RBCs.
السيت أعلى
- * It can also reflect less plasma.
الحقن أقل

- * Clinical interpretation must distinguish the two.

Since each indicate a specific condition.

- * increased hematocrit due to ↓ plasma ⇒ Hypovolemia (Dehydration)
- * increased hematocrit due to ↑ RBCs ⇒ polycythemia.

Hematology

RBC



we reached Here.

Red Blood Cells/ Erythrocytes

