




MIND MAPPING


Histology mind maps for midterm
Done by:- TOLEEN MAWAZREH

LECTURE 1


Blood as a connective tissue



Blood




Functions of Blood




HEMATOCRIT: Ratio of the volume of RBCs to the volume of whole blood
 Example: a hematocrit value of 40% means that there are 40 ml of RBCs in 100 ml of whole blood
 Normal hematocrit:
 Males=40-53%
 Females= 36-48%

Fluid




Preparing a blood smear




Blood cells can be studied histologically in smears prepared by spreading a drop of blood in a thin layer on a microscope slide

Polychromatic stains



Collected blood in which clotting is prevented by the addition of anticoagulants (eg, heparin, citrate or EDTA) can be separated by centrifugation into layers that reflect its heterogeneity



LECTURE 2

Erythropoiesis

Production of erythrocytes in the bone marrow, is stimulated by erythropoietin



Erythropoietin is produced by the kidneys



When RBC count drops, such as during blood loss, the resulting oxygen-deficiency state, hypoxemia, is detected by the kidneys.



The kidneys respond by increasing their erythropoietin secretion, which leads to increased red blood cell production

Polycythemia: an increase in the total number of RBCs (and/or hemoglobin)
While Anemia: is a decrease

People living at high altitudes usually have higher RBC count as a response to lower oxygen levels.

Athletes whose demand for oxygen is more elevated, also have higher RBC counts.


	RBCs (Erythrocytes)	WBCs (Leukocytes)
Types	1 type	5 types
	Not true cells	True cells
Number	Male: 4.5-5.5 million/mm3 female:4-5 million/mm3	4500-11000/mm3
Diameter	6-9um (7.5um)	6-20um
Life span	120 days	Few days-years
Origin and maturation	Bone marrow	Bone marrow and lymphoid tissue
Shape	Biconcave discs	Spherical
Function	Gas exchange	Defense
Motility	Non motile	Motile
	Function exclusively within vascular system	Function mainly OUTSIDE blood vessels in the tissues

Leukocytes

- Originate in the bone marrow and released continuously into the blood
- **Travel** in bloodstream but **function** mainly outside blood vessels (in loose CT)
- Leukocytes form a mobile army that helps protect the body from damage by bacteria, viruses, parasites, toxins and tumor cells

Thrombocytes (Platelets) neither granulocytes nor agranulocytes

- Small non-nucleated cytoplasmic fragments
- Formed by fragmentation of the cytoplasm of megakaryocytes in the bone marrow
- Number: 200,000-400,000/mm3
- Shape: biconvex discs
- Cytoplasm: purple, granular
- Diameter: 2-4 um
- Lifespan about 10 days
- Control the bleeding by plugging the defects in blood vessels and activating blood clotting cascades



In stained blood smears, platelets often appear in clumps

Platelete has 2 Zones

Outer pale basophilic (clear) peripheral zone: hyalomere

Hyalomere: contains cytoskeleton and membranous channels

- Cytoskeletal elements

> Microtubule

> Actin filaments

Maintain shape and help contractions of platelets and squeezing, clot retraction

Membrane channels

→ Open canicular system

→ Dense tubular system

Central dark granular zone: granulomere

Granulomere: contains granules and organelles

Alpha granules: clotting factors, growth factors

Dense (delta) granules: serotonin (absorbed from plasma), ATP, ADP

Lambda granules: lysosomes (aid in clot resorption)



Have thick glycocalyx

Microtubule and microfilament

Open canicular system

Dense tubular system

H.K.

Delta granule

Alpha granule

Lambda granule

Their main function is to continually monitor the vascular system and detect any damage to the endothelial lining of the vessels. If the endothelial lining breaks, the platelets adhere to the damaged site and initiate a highly complex chemical process that produces a blood clot Thus preventing blood loss

Specific granules

Secondary granules

Cytoplasmic granules

Non-specific granules

Azurophilic granules

Lysosomes

Primary granules

Classification


Depending on the presence or absence of specific granules in their cytoplasm

Granulocytes

- Cytoplasmic granules (containing enzymes or chemicals) → makes cytoplasm look grainy
- Single multi-lobed nucleus (segmented)
- All are phagocytic; they engulf and consume foreign cells and material
- 3 main types:

Neutrophils

Small granules, pale pink/ salmon pink



Neutrophil

Eosinophil


Large granules, Red



Eosinophil

Basophils

Large granules, blue



Basophil

Agranulocytes


Monocytes

Lymphocytes

	Neutrophil	Eosinophil	Basophil
Percentage (WBCs)	Most		Least
Size	12-15um	12-15um	12-15um
Life span	Few days	Few days	Few days
Nucleus	2-5 lobes	2 lobes	Irregular (S-shaped)
Phagocytic activity	Most		Least
Motility	Most		Least




Never Let Monkeys Eat Bananas
Most common to least

- Rarest leukocyte – might not see these under the microscopes
- Usually have bi-lobed, S-shaped nuclei obscured by the large basophilic granules
- Has large granules that stain dark purple/ blue in basic dyes (basophil = basic loving)
- Granules contain histamine, heparin and eosinophilic chemotactic factor that mediate inflammation in **allergic reactions and parasitic infections**



- The most common leukocyte
- 2-5 lobes in nucleus connected by “threads” of nuclear material (polymorphs)
- Light pink cytoplasm
- Called neutrophils because cytoplasm takes up red (acidic) and blue (basic) stains equally
- Specialized for responding to **Bacterial invasions- Acute infections- Acute pyrogenic infections**
- Neutrophils are short-lived cells with a half-life of 6-8 **hours in blood** and a life span of 1-4 **days in connective tissues** before dying by apoptosis

Note: Neutrophils will show increasing segmentation (many segments of the nucleus) as they mature. A normal neutrophil should have 3–5 segments. Hypersegmentation is not normal but occurs in some disorders, most notably vitamin B12 deficiency.



Different names for neutrophils:

Polymorphs / Pus cells

Myelocytes / Microphages

Cells of acute inflammation


Basophils

Eosinophils

Granulocytes

Neutrophils

- Usually have bi-lobed nuclei connected by a short “thread” of nuclear material
- Large cytoplasmic granules, which stain red with the acidic eosin dye (eosinophil = eosin loving)
- Help in **ending allergic reactions and in fighting parasitic infections**




General features of granulocytes

- Spherical in blood stream, irregular in connective tissue
- Highly motile with different shapes due to their amoeboid movement
- Leave blood stream by migrating between the endothelial cells by a process called diapedesis


Oval in shape, with flattened crystalloid cores

Externum (pale):
contains histaminase and sulfatase



Internum (dark):
contains basic protein to kill parasites

Specific granules (Crystalloid granules)



Pyogenic is referring to bacterial infections that make pus while **pyrogenic** is producing heat

The ways of killing

Oxygen-dependent killing:


The neutrophil undergoes a ‘respiratory burst’ activating NADPH oxidase to produce reactive oxygen species. Hydrogen peroxide is formed, and with the help of myeloperoxidase, it generates hypochlorous acid.

Oxygen-independent killing:

Granules inside the neutrophil release enzymes like defensins, lysozyme, and proteases. These proteins break down microbial walls and deprive bacteria of nutrients.


NETs

the Neutrophil Extracellular Trap, or NET. When overwhelmed, a neutrophil can undergo NETosis—a unique cell death where it releases its DNA coated with antimicrobial proteins. This sticky web traps and kills microbes outside the cell, preventing them from spreading.”



Barr body

In females, the inactive X chromosome (Barr body) may appear as a drumstick-like appendage on one of the lobes of the nucleus (about 3% of neutrophils in peripheral blood)



Cells of acute infection


Neutrophils are the first WBCs that leave the blood in large numbers to reach the site of inflammation

Why???

- 1- The most abundant
- 2-The most motile
- 3- Neutrophil chemotactic factors are the first released

PUS

Dead neutrophils, bacteria, lysed ECM, and tissue-fluid form a viscous, usually yellow collection of fluid called pus.



Agranulocytes

- Single non-lobulated nucleus
- Granules in cytoplasm are too small to see (**nonspecific granules**, azurophilic granules, primary granules, lysosomes)

Monocytes

- Largest leukocytes
- Bluish cytoplasm (frosted glass appearance) & a large C-shaped nucleus
- Highly motile and phagocytic
- Travel through bloodstream to reach connective tissues, where they transform into macrophages (large phagocytic cells)

Perform 2 functions → phagocytosis and antigen presentation

Are precursor cells of macrophages, osteoclasts, microglia, and other cells of the **mononuclear phagocyte system** in connective tissue

All monocyte-derived cells are **antigen-presenting cells**

all characterized by phagocytic activity

Kupffer Cells
Liver

Microglia
CNS

Dust cells
Lung

Osteoclasts
Bone resorption

Langerhans cells
Epidermis

Dendritic cells
Lymph node
Spleen


Macrophages
Bone marrow
Connective tissue

Monocytes and macrophages are the same cells at different stages of maturation



BONE MARROW **BLOOD** **TISSUE**



Morphologic changes occur upon differentiation of monocytes into macrophages:




- The cell increases in diameter more than threefold
- The cell acquires more cytoplasmic granules and vacuoles
- Cell surface receptors increase in number
- The phagocytic activity improves



B Lymphocyte

(Humoral immunity)

- Differentiate into plasma cells
- Secrete antibodies that bind to specific antigens and mark them for destruction by phagocytic cells





Produces antibodies

Memory cells allow a more rapid response if the antigen appears again later which allows a very rapid response upon subsequent exposure to the same antigen. Basis of immunity/vaccination

Immunohistochemistry

Direct method



Lymphocytes

Morphologically lymphocytes can be classified into

Large
(9-18 micro m)
Active lymphocyte

Small
(6-9 micro m)
Inactive lymphocyte

The amount of cytoplasm depends upon state of activity of the lymphocyte
In circulation blood there is predominance of small inactive lymphocytes

Could be

B lymphocyte
T helper
T cytotoxic
Natural killers

Could be

B lymphocyte
T helper
T cytotoxic

- Smallest leukocytes
- Round nucleus occupies most of cell volume
- Cytoplasm is light clear blue
- Increased numbers are commonly seen in **viral infections**
- Lymphocytes vary in life span according to their specific function, some live for a few days and some live for many years

T Lymphocyte

(Cell mediated immunity)

Has different types, some directly kill foreign or infected cells; others activate phagocytes to destroy microbes

T helper (CD4+) **T cytotoxic (CD8+)** **Suppressor T cell (CD8+ or CD4+)**

Help cytotoxic T cells and B cells in their immune functions

Kill virus-infected, transplanted and neoplastic cells (**adaptive immunity**)


Suppresses immune response to self Ag
Suppresses immune response of T and B lymphocytes

The retrovirus that produces acquired immunodeficiency syndrome (AIDS) infects and rapidly kills helper T cells. Reduction of this key lymphocyte group cripples the patient's immune system rendering them susceptible to opportunistic bacterial, fungal, protozoan, and other infections that usually dealt with easily in immunocompetent individuals.

Suppressor T cells switch off the immune response when the stimulus is removed
Damage to suppressor cells can result in autoimmune disease

Natural killer cells (NULL Lymphocyte)




Kill virus-infected, transplanted and neoplastic cells (**innate immunity**)



Large granular lymphocytes
Activated lymphocytes


Similar to cytotoxic T cell
But T cell needs activation by MHC molecules while NKC are always large and activated

Natural killer cells don't have a problem with MHC1 molecules, but once it sees a cell without those MHCs it will kill it directly



The antigens presented by class I peptides are derived from intracellular proteins (cytosolic)



Major histocompatibility complex MHC










T lymphocytes are specialized to recognize both classes of MHC proteins and the antigens they present

If the MHCs on cells of a tissue graft are not similar to those that T lymphocytes encountered during their development, the grafted cells will induce a strong immune reaction by T cells of the recipient.

To these lymphocytes, the unfamiliar MHC epitopes on the graft's cells are recognized as markers of "non-self" cells that they must eliminate.











Between the hematopoietic cells run the **sinusoids**, which have discontinuous endothelium, through which newly differentiated blood cells and platelets enter the circulation

Red marrow is also a site where older, defective erythrocytes undergo phagocytosis by macrophages, which then reprocess heme-bound iron for delivery to the differentiating erythrocytes.



Blood Cell Formation (Hematopoiesis)

Erythropoiesis

the process which produces erythrocytes

Granulopoiesis

the process which produces granulocytes

Thrombopoiesis

the process which produces thrombocytes

Lymphopoiesis

the process which produces lymphocytes

Monocytopoiesis

the process which produces monocytes

Distribution of RBCs and WBCs in

peripheral blood

RBCs → in 1 ul of peripheral blood → 5 million/ul

WBCs → in 1 ul of peripheral blood → 4500-11000/ul


bone marrow

Myeloid: Erythroid 3:1




Erythropoiesis (red cell formation)

- Takes about 1 week
- Rate is controlled by the hormone erythropoietin (secreted by the kidney cells) and the availability of iron, folic acid, vitamin B12, protein precursors

- Normally, only about 1% of all red blood cells in the bloodstream are reticulocytes
- They circulate for about 1-2 days before developing into mature red blood cells
- An increase in reticulocytes ---- blood loss (hemorrhage)




Reticulocyte Mature Erythrocyte



Why?! As it should be the opposite?

Due to the life span, WBCs die within 2-5 days after eliminating infection, so there should be compensation of this loss, so bone marrow produce them in larger numbers



White blood cell abnormalities

In peripheral blood

Increased numbers of white cells appear in the peripheral blood in a variety of disorders and provide a useful clue to the underlying disease.

considerable and sustained increase of **circulating neutrophils** in bacterial infection

An increase of **circulating eosinophils** in parasitic infection and some allergies

In bone marrow

Suppression of these cells on bone marrow

Hypoplastic bone marrow
Aplastic anemia (pancytopenia)

Malignant proliferation of these cells inside bone marrow

Leukaemia

Vast number of white cells and their precursors (many of which spill over into the blood)
Leukemia is classified according to the cell line involved (granulocytic, monocytic, lymphocytic)


Clinical applications

Bone marrow Aspirate or biopsy

Needed to diagnose disorders like aplastic anemia or leukemia

Bone marrow transplantation

In bone marrow diseases like leukemia, hematopoietic stem cells taken from a donor are infused into the same or another person



Stage	Cell size مقياس مطلوب	Nucleus	Cytoplasm	Notes
Proerythroblast	Largest cell (17um)	Large pale nucleus with prominent nucleolus	Pale basophilic cytoplasm	
Basophilic erythroblast	The cell becomes smaller (15um)	smaller and darker	Deeply basophilic cytoplasm (high in ribosomes)	
Polychromatophilic erythroblast	The cell becomes smaller (13um)	smaller and darker	Cytoplasm becomes violet	Takes basic (ribosomes) and acidic stains (Hb)
Normoblast Orthochromatic erythroblast	The cell becomes smaller (11um)	smaller, darker and eccentric to be expelled outside	Cytoplasm is acidophilic (Hb)	The nucleus is extruded at this stage
Reticulocyte	Immature erythrocyte but slightly larger (9um)	No nucleus	Cytoplasm is acidophilic (Hb) but contains remnants of ribosomes forming reticulum	Can be stained by supravital stains (brilliant cresyl blue)
Erythrocyte				

Stages of differentiation are characterized by:
1- Decreasing cell size
2- Progressive loss of organelles


Presence of free ribosomes at early stages

Accounts for the marked cytoplasmic basophilia (**blue**)



3- Progressive increase in hemoglobin content

Accounts for increasing eosinophilia (**pink/red**)



Blood Cell Formation (Hematopoiesis)



Granulopoiesis (Neutrophils, Eosinophils and Basophils formation)




Neutrophilic band cells (important CLINICALLY)
Their percentage does not exceed 5% in peripheral blood
The appearance of large numbers of immature neutrophils (band cells) in the blood, sometimes called a "shift to the left," is clinically significant, usually indicating a **bacterial infection**.
This means that the bone marrow has been signaled to release more neutrophils and increase production of neutrophils



Changes in the number of neutrophils in the blood must be evaluated by taking all their compartments into consideration.



Pools of neutrophils

four functionally and anatomically defined compartments for Developing and maturation of neutrophils



Neutrophilia

an increase in the number of **circulating neutrophils**




LECTURE 5

Lymphatic system

Components

- **lymphatic fluid**
Lymph is a colorless fluid that floats in the lymphatic vessels. It is similar in composition to blood plasma
- **lymphatic vessels**
thin vessels that accompany arteries and veins throughout the body and transport lymph.
- **lymphatic tissue**
specialized form of reticular connective tissue that is composed of masses of lymphocytes.
- **lymphatic organs**
include the lymph nodes, spleen, thymus, and red bone marrow

- occur alone as lymph nodules (follicles)
- organized into various lymphatic organs.




functions

- Drain excess interstitial fluid from the tissues and return to blood stream
- Initiate an immune response against disease by producing and transporting lymphocytes
- Transport dietary lipids absorbed by the gastrointestinal tract into the blood

Some lipids are too large to pass through the capillary walls of the small intestine and therefore cannot be absorbed.

lymphatic system is used, for example, for dietary fat absorption in the intestine



The lymphatic capillaries within the small intestine, known as **lacteals**, can absorb these large lipid molecules and transport them into the venous circulation via the thoracic duct. Lymph containing these lipids becomes a creamy white color and is referred to as **chyle**.

Lymphatic Organs


Lymphocytes can be found throughout the body, however, they **aggregate in places** where they are most likely to come into contact with pathogens.

Primary lymphatic organs

- Bone marrow.
- Thymus gland.

Are sites of Lymphocyte production, maturation,

selection



Secondary lymphatic organs


- Diffuse lymphatic tissue (lymphatic nodules).
- Spleen.
- Lymph nodes.

Are sites to encounter pathogens and become activated

Lymph nodes


Consists of

- Reticular tissue forms the stroma of the lymph node
- Immunocompetent B cells and T cells are suspended throughout the lymph node



Functions


- Nodes filter the lymph, removing foreign material and microorganisms.
- Antibody-mediated and cell-mediated immune responses occur in the lymph nodes




Lymph node enlargement

- in cases of lymphoma (painless lymphadenopathy)
- infection (painful)

Lymphatic capillaries are made of overlapping endothelial cells with **large gaps**. The overlapping flaps function as a **one-way valve**.



Around 20 liters of fluid leaves the arterial capillaries every day, but only 17 liters of fluid returns to the venous capillaries. Lymphatic vessels function to drain this excess fluid (3L) from the tissues as lymph and return this fluid to the blood.



Lymph returns back to the big veins (venous angle: the junction between subclavian and internal jugular veins) through the Thoracic duct and Right lymphatic duct.

Lymph flow is enabled by the same forces that facilitate blood flow in the veins. It goes from lymphatic capillaries to larger and larger lymphatic vessels and eventually drains into the bloodstream. On the way, it passes through a number of lymph nodes, which serve as filters, cleansing the fluid before it reaches the bloodstream.

When fluid accumulates in the tissue

- interstitial pressure increases pushing the flaps inward
- opening the gaps between cells, allowing fluid to flow in
- pressure inside the capillary increases
- the endothelial cells are pressed outward, closing the gaps, thus preventing backflow.

Unlike blood capillaries, the gaps in lymphatic capillaries are so large that they allow bacteria and immune cells (ex. Macrophages/ dendritic cells) to enter. This makes the lymphatic system a useful way for large particles to reach the bloodstream.

Lymph nodes

covered by


consists of

outer cortex

- highly cellular part of the lymph node
- Contains lymphatic follicles
- Receives lymph from **afferent vessels**

inner paracortex
contains mostly T-cells

outer cortex
has lymphatic follicles that mostly contain B-cells



Other cells in the lymph node:
Macrophages
Dendritic cells
Follicular dendritic cells
Reticular cells

inner medulla


- No follicles
- Forms sinuses that lead to **efferent vessels** at the hilum

contain mostly **plasma cells**

Lymphocytes


Small
(6-9 micro m)

Inactive lymphocyte
Darkly stained cell




Large
(9-18 micro m)

Active lymphocyte
Lightly stained cell



Note: the nucleus of plasma cell is eccentrically placed, round or oval with *cartwheel* (clock-face) chromatin pattern.



composed of

medullary cords


Separated by

medullary sinuses

converge at

the hilum

slight indentation on one side of the node. Here, an artery, vein, and an efferent lymphatic vessel enter and leave the node.



a **capsule** of dense connective tissue, and have capsular extensions called the **trabeculae**, which provide support for blood vessels entering into the nodes. When lymph nodes become enlarged, the capsule is stretched and becomes painful

Other cells in the lymph node

Macrophages

Dendritic cells

Follicular dendritic cells

Reticular cells

Monocytes derived cells


	Macrophage	Dendritic cell	Follicular dendritic cell
Phagocytosis	Most phagocytic	Moderately phagocytic	X
Antigen presenting (via MHC-II)	Moderate Ag-presenter	Very powerful Ag-presenter	X
Location in lymph node	Cortex and medulla	Cortex and medulla	Outer cortex

Are antigen **HOLDING** cells
Holds the Ag for long time

Macrophages and Dendritic cells capture antigen within tissues and transport antigen to secondary lymphoid tissue

Lymphatic trunks and ducts

All lymphatic vessels coalesce to form larger trunks which eventually converge to form the right lymphatic duct and the thoracic duct



Right lymphatic duct

- Is formed by right jugular + + right subclavian trunks
- Drains lymph from the upper right quadrant of the body (the right side of the head and neck, the right side of the thorax and the right upper limb)
- Empties into the junction where right internal jugular vein joins the right subclavian vein (**Rt venous angle**)

Thoracic duct (Left lymphatic duct)

- Is larger and drains lymph from the rest of the body
- Originates in the abdomen as **cisterna chyli**
- Passes through the diaphragm at the aortic aperture
- Empties into the junction where left internal jugular vein joins the left subclavian vein (**Lt venous angle**)


Cisterna chyli is a dilated sac at the lower end of the thoracic duct (anterior to the bodies of L1 and L2) formed by confluence of the right and left lumbar trunks and the intestinal trunk

Efferent vessels

The lymph is removed from the medullary sinus via one or two efferent lymphatic vessels that leave the lymph node at the hilum. Valves in the vessels prevent lymph from flowing in the wrong direction.

Efferent lymph contains lots of activated T-lymphocytes, activated B-lymphocytes, plasma cells and antibodies.

Sinuses are irregular spaces through which the lymph percolates
All the lymphatic sinuses are lined by a discontinuous layer of simple squamous endothelium



Lymph, containing **micro-organisms, soluble antigens and antigen presenting cells**

Afferent vessels

Many afferent lymphatic vessels enter the lymph node at different points over its convex surface, each containing valves to prevent backflow of lymph.

Subcapsular sinuses

Each afferent vessel empties into the subcapsular sinus

Trabecular sinuses

a continuation of the subcapsular sinuses that follow the trabeculae and drain into the medullary sinuses.

Medullary sinuses

Found separating the cords. The medullary sinuses converge at the hilum into the efferent lymphatic vessel.

Lymph slows down when it passes in lymph nodes.


Lymphocytes can enter lymphoid tissues in two ways

consists of

Direct entry into lymph nodes via afferent lymphatics

Entry from blood capillaries across specialized endothelial cells present in the postcapillary venules (High Endothelial Venules= HEV) within the paracortex of the lymph node


Primary follicles
lymphoid follicles without a germinal center. (virgin B cells)



As B cells in lymphatic follicle are stimulated, they differentiate into plasma cells. Plasma cells move to medulla (medullary cords)

Secondary follicles
lymphoid follicles with a germinal center. Sites for B memory cell and plasma cell generation

When activated by antigens (and T helper cells),
B cells migrate to the center of the follicle, forming a germinal center. Germinal centers are the central regions of secondary follicles where activated B cells are proliferating (dividing by mitosis) and differentiating into plasma cells and memory B cells. When stimulated by antigens, lymph nodes enlarge due to the formation of germinal centers and B cell proliferation



LECTURE 6

Spleen

- ✓ an oval-shaped intraperitoneal organ
- ✓ Has a notched anterior border.
- ✓ lies high on the upper left portion of the abdomen, just beneath the diaphragm, behind the stomach and above the left kidney
- ✓ the **largest of the lymphoid organs**

Functions


- **Filtration of blood** (defense against blood-borne antigens)
- The main site of **old RBCs destruction**.
- **Production site of antibodies and activated lymphocytes** (which are delivered directly into the blood)

Spleen

The splenic artery

the largest branch of the celiac artery. It has a tortuous course as it runs along the upper border of the pancreas. The splenic artery then divides into about six branches, which enter the spleen at the hilum

supplies the spleen as well as large parts of the stomach and pancreas



The splenic vein

leaves the hilum and runs behind the tail and the body of the pancreas. Behind the neck of the pancreas, the splenic vein joins the superior mesenteric vein to form the portal vein

In cases of portal hypertension, spleen often enlarges from venous congestion.


The spleen is covered by a **capsule** of dense connective tissue

have

capsular extensions called the **trabeculae**

have

Large trabeculae **originate at the hilum**, on the medial surface of the spleen, and carry branches of the splenic artery, vein, lymphatics, and nerves into the spleen



Blood flow through the spleen

Each central arteriole eventually leaves the white pulp and enters the red pulp, losing its sheath of lymphocytes and branching


penicillar arterioles

several short straight penicillar arterioles that continue as terminal capillaries.

Terminal capillaries

(Sheathed capillaries)

Some of these terminal capillaries are sheathed with APCs for additional immune surveillance of blood



Splenic sinusoids

Trabecular veins



Splenic vein

Spleen

Stroma

Reticular tissue (reticular fibers and reticular cells)

parenchyma



Marginal zone sinuses


Red pulp

which appears red on gross examination (**blood filled**)

White pulp

lymphoid tissue

contains



When the lymphatic sheath expands to incorporate the follicles, the central arteriole is displaced to one side and acquires an eccentric position in the follicle but is still called the central arteriole (Follicular arteriole).

Lymphoid follicles

spherical aggregations of **B cells** scattered throughout the PALS

These follicles have the same structural organization as those found in lymph nodes

→ **Primary (unstimulated) follicles** contain resting (inactive) B cells

→ **Secondary (stimulated) follicles** contain activated B cells in a central region (germinal center)

Production of antibodies and activated lymphocytes (which are delivered directly into the blood)

Splenic nodules (Malpighian corpuscles)

Constituting 75% of the spleen, the red pulp is responsible for the hematological (circulatory) function of the spleen.

contains

Splenic cords (Billroth's cords): consist of all cells between the sinusoids in the red pulp (reticular cells, macrophages, plasma cells, lymphocytes, RBCs, platelets, other leukocytes)

Splenic sinusoids: blood-filled spaces located throughout the red pulp. They have large, dilated, irregular lumens and large pores (spaces between the endothelial cells)



Function: Destruction of worn-out RBCs and platelets

which appears white on gross examination (**collection of both B and T lymphocytes**)

Constituting 25% of the spleen, the white pulp is responsible for the immunological (lymphatic) function of the spleen.


Periarteriolar lymphatic sheaths (PALS)

tightly packed **T cells** arranged in cylindrical sheaths around central arterioles




Function: The lymphocytes and APCs monitor the blood for foreign antigens and respond in a similar way to those in the lymph nodes.


Red pulp



When B cells in the primary follicles are exposed to Antigen, they proliferate and differentiate to plasma cells and move toward the red pulp.



wide gaps between endothelial cells which allow for movement of entire cells from cords => to sinuses




- 1. The endothelial cells (stave cells) are elongated, fusiform cells that lie parallel to the long axis of the vessel
- 2. The cells lie side by side around the vessel but not joined by any type of intercellular junctions
- 3. The endothelial cells are supported by highly discontinuous basal lamina (forms bars and encircles the sinusoid)

After surgical removal of the spleen (splenectomy), the number of abnormal erythrocytes in the circulation increases although most such cells are then removed by macrophages in sinusoids of the bone marrow and liver.

Blood flow through the splenic red pulp can take either of two routes:


- Open circulation: the capillaries open into the spaces of the red pulp (splenic cords) and then the blood returns to the venous system through the wall of the splenic sinusoids
- Closed circulation: the capillaries open directly into the splenic sinusoids (blood is enclosed by endothelium)



Marginal zone sinuses

- Located between the white and the red pulp
- The spaces between these sinuses are wide (2-3um)

It is here the blood- borne antigens and particulate matter have their first free access to the parenchyma of the spleen



events occur at the marginal zone:



- 1- APCs sample the material travelling in blood searching for antigens
- 2- Macrophages attack microorganisms present in the blood
- Lymphocytes come into contact with APCs, if they recognize their Ag-MHC complex, the lymphocytes initiate immune response within the white pulp
- 3- The circulating B and T cells leave the blood stream to enter the preferred location within the white pulp
 - T cells: PALS
 - B cells: lymphatic follicles

Functions of the spleen

- circulatory functions
 - Blood cell production: During the fetal life, blood cells are produced in the spleen
 - Blood storage: A small quantity of blood is stored in the sinusoids of the red pulp
 - RBC destruction: Most worn-out or damaged red blood cells are destroyed in the spleen (some in the liver and bone marrow). They are phagocytized by macrophages
- lymphatic functions
 - Defense mechanism
 - Macrophages phagocytize microbes that have penetrated the blood.
 - Antigens in the blood activate B and T cells residing in the spleen, triggering immune response
 - Production of antibodies and activated lymphocytes (which are delivered directly into the blood)

Lymph node vs spleen

Lymph node	Spleen
Multiple, small	Single, large
Along the course of lymphatic vessels	Intra-abdominal
Filters lymph	Filters blood
Covered by fascia	Covered by peritoneum
Has afferent vessels	No afferent vessels
Cortex and medulla	White pulp and red pulp
Contains Lymphatic sinuses	Contains Blood sinuses



LECTURE 7


Diffuse lymphatic tissue (lymphatic nodules)

- Is formed by aggregations of lymphatic tissue
- Is found in various **mucosal sites of the body**


- inner lining of the
- digestive tract
 - respiratory tract
 - genitourinary tract

a common site of invasion by pathogens because their lumens open to the external environment.

therefore be referred to as









Mucosa-Associated Lymphatic Tissue (MALT)




Tonsils

are large, irregular masses of lymphoid tissue
Function of tonsils: Protect the body from inhaled and ingested pathogens.




Adenoids



With the surface epithelial cells removed, scanning electron microscopy (SEM) shows typical **basement membrane over the villi (V)** but reveals a highly porous covering over **lymphoid nodules** of the Peyer patch.


This sieve-like basement membrane facilitates interactions between immune cells and M cells in the epithelium over the nodules

Gut-associated lymphoid tissue (GALT)




Examples

Peyer's patches of ileum



Lymphatic nodules of appendix




Peyer's patch and M cells

antigens in the gut lumen are bound by **M cells** and undergo transcytosis into their intraepithelial pockets

dendritic cells take up the antigen, process it, and present it to **T helper cells**


B lymphocytes stimulated by the **Th cells** differentiate into **plasma cells** secreting IgA antibodies

The IgA is transported into the gut lumen where it binds its antigen on the surface of microorganisms, neutralizing potentially harmful invaders before they penetrate the mucosa.



Bronchus-associated lymphoid tissue (GALT)

- Is located in the mucosa of the bronchioles.
- Function: Protects the body from inhaled pathogens.



LECTURE 8

Thymus

✓ The thymus is a bi-lobed gland located in the anterior mediastinum, posterior to the sternum and anterior to the trachea.

✓ It is **large in the newborn** and young child


From puberty onwards, it gradually **becomes replaced by fat**

Fully formed and functional at birth, the thymus remains large and very active in T-cell production until puberty during which it normally undergoes involution, with **decreasing lymphoid tissue mass and cellularity** and reduced T cell output => may be involved with the decline of immune function in the elderly

Within the thymus

- **immature T-cells**
 - develop
 - differentiate
 - multiply
- **gaining their antigen specificity and immune tolerance to the body's own tissues**


Positive selection



selection

Only about 2% of all developing T lymphocytes pass both the positive and negative selection tests and survive to exit the thymus as **immunocompetent T cells**.

negative selection



Double function

As gland part of the endocrine system

- Secreting thymic **hormones**
- Thymosin
 - Thymopoietin
 - Thymotaxin

As a primary lymphatic organ

For production and maturation of T-lymphocytes


Double embryonic origin

Endoderm

Originates from the embryo's **third pair of pharyngeal pouches**

↓

unique thymic epithelial cell



Fully mature T cells (immunocompetent T-cells) leave the medulla via venules and efferent lymphatic vessels

- nodes (paracortex)
- the spleen
- diffuse lymphatic tissues

they reside and are responsible for **cell-mediated immune responses**

After maturation in primary lymphoid organs

↓ B and T cells circulate to

the peripheral secondary lymphoid organs (the MALT, the lymph nodes, and the spleen)

Lymphocytes do not stay long in the lymphoid organs; they continuously recirculate through the body in connective tissues, blood, and lymph.

If no antigen is present

lymphocytes routinely enter and leave secondary lymphoid tissues

If antigen enters the secondary lymphoid tissue



Lymphocyte proliferation in response to antigen occurs within the lymphoid tissue.

After several days, antigen-activated lymphocytes begin leaving the lymphoid tissue.

Mesoderm

Hematopoietic origin

Immature T lymphocytes (T lymphoblasts) circulating from the bone marrow to invade and proliferate in thymus during its development.



The staining differences reflect the much greater density of lymphocytes in the cortex than the medulla

outer darkly basophilic cortex

- The cortex contains
- Immature T cells (T lymphoblasts, thymocytes) (in various stages of differentiation and maturation)
 - Macrophages

As T cells mature, they migrate to the medulla

more lightly stained medulla

- The medulla contains
- Fewer and more **mature** lymphocytes.
 - Macrophages
 - Dendritic cells (APCs)
 - Unique thymic epithelial cells (TECs)
 - Large aggregates of TECs called **Hassall corpuscles**

- are unique to the thymic medulla
- Up to 100 µm in diameter
- Are concentric aggregates of squamous cells with central keratinization (**acidophilic**)
- Tend to grow larger with age




Unique thymic epithelial cells (TECs) **Develop from endoderm**


Epithelial reticular cells

- Form a **stroma** to which macrophages and developing lymphocytes attach instead of reticular fibers
- Envelop groups of T cells that are multiplying and maturing (in cortex)
- Act as APCs, expressing MHC class II and MHC class I molecules (in cortex)
- Express many specialized proteins specific to cells of other organs, tissue specific antigens (in medulla)
- Secrete hormones that promote the differentiation of T cells (endocrine thymus)
- Line the capsule and septa and surround all blood vessels in the cortex

Form a **blood-thymus barrier** preventing antigens in the blood from making contact with the developing T cells (in cortex)



called **Nursing Cells**



Advantages of lymphocyte recirculation

- Lymphocyte recirculation enables the limited number of naive lymphocytes in an individual that are specific for a particular antigen to search for that antigen throughout the body
- It ensures that particular lymphocytes are delivered to particular tissue

Recirculation of naive lymphocytes: recirculate through secondary lymphoid organs

Recirculation of activated lymphocytes: migrate to peripheral tissues at sites of infection

Ex.

If you have a viral infection in your oral cavity, this infection spreads to lymph nodes draining the oral cavity, and lymphocytes in these lymph nodes develop an immune response against this virus.

Then the activated lymphocytes leave the lymph node and circulate in the lymph and then blood.

They will go to every lymph node in your body and settle there

↓

The whole body is prepared against this virus., the whole body is tuned up for the next infection with the same virus