
Heart Pump and Cardiac Cycle


Faisal I. Mohammed, MD, PhD

Yanal A. Shafagoj, MD, PhD

Objectives

- To understand the volume, mechanical, pressure and electrical changes during the cardiac cycle
- To understand the inter-relationship between all these changes
- To describe the factors that regulate Cardiac output and Stroke volume.
- Resources: Textbook of Medical Physiology By Guyton and Hall


(a) ECG
R
T
P
Q
S
T
(b) PRESSURE (mm Hg)
120
100
80
60
40
20
0
Aortic semilunar valve closes
Dicrotic wave
Bicuspid valve closes
Aortic semilunar valve opens
Bicuspid valve opens
Aortic semilunar valve closes
Dicrotic wave
Bicuspid valve opens
Aortic pressure
Left ventricular pressure
Left atrial pressure
(c) VOLUME OF LEFT VENTRICLE (ml)
130
60
0
(d) HEART SOUNDS
S1
S2
0.4 sec.
0.1 sec
0.3 sec
Relaxation (quiescent) period
Atrial systole
Ventricular systole
1
Isovolumetric relaxation
2
Rapid
Diastasis
Atrial systole
Ventricular filling
3a
Isovolumetric contraction
3b
Ventricular ejection
Isovolumetric relaxation
1
Ventricular filling
Atrial systole
2
Isovolumetric contraction
Ventricular ejection
3
(e) PHASES OF THE CARDIAC CYCLE


Time (seconds)
0
0.3
0.6
0.9
Pressure (mm Hg)
120
100
80
60
40
20
0
Aortic valve opens
Aortic valve closes
A-V valve closes
A-V valve opens
Aortic pressure
Atrial pressure
Ventricular pressure
Volume (ml)
160
120
80
Ventricular volume
R
P
Q
S
T
Lub
Dup
Electrocardiogram
Heart sounds
Ventricular systole
Ventricular diastole
Ventricular systole

Cardiac Cycle

- Cardiac cycle refers to all events associated with blood flow through the heart
  - Systole – contraction of heart muscle
  - Diastole – relaxation of heart muscle

Cardiac Cycle

- Atrial systole 0.1 second
- Atrial diastole 0.7 second
- Ventricular systole 0.3 second
  - Isovolumic contraction 0.01 seconds
  - Rapid ejection period
  - Slow ejection period
- Ventricular diastole 0.5 seconds
  - Isovolumic relaxation 0.02 seconds
  - Rapid filling
  - Slow filling (Diastasis)
  - Atrial contraction

Cardiac cycle …cont

- End diastolic volume (EDV) – End systolic volume (ESV) = Stroke volume (SV)
- SV X heart rate (HR) = cardiac output (CO)
- Ejection fraction = SV/EDV
- Inotropic vs. Chronotropic
- Autonomic control of cardiac cycle (pump)

Phases of the Cardiac Cycle


- Ventricular filling – mid-to-late diastole
  - Heart blood pressure is low as blood enters atria and flows into ventricles
  - AV valves are open, then atrial systole occurs

Phases of the Cardiac Cycle


- Ventricular systole
  - Atria relax
  - Rising ventricular pressure results in closing of AV valves
  - Isovolumetric contraction phase
  - Ventricular ejection phase opens semilunar valves

Phases of the Cardiac Cycle

- Isovolumetric relaxation – early diastole
  - Ventricles relax
  - Backflow of blood in aorta and pulmonary trunk closes semilunar valves
- Dicrotic notch – brief rise in aortic pressure caused by backflow of blood rebounding off semilunar valves


Left heart
Electrocardiogram
Heart sounds
QRS
P
T
P
1st
2nd
Pressure (mm Hg)
120
80
40
0
Dicrotic notch
Aorta
Atrial systole
Left ventricle
Left atrium
Ventricular volume (ml)
130
100
70
EDV
SV
ESV
Atrioventricular valves
Aortic and pulmonary valves
Phase
Open
Closed
Open
Closed
Open
Closed
1
2a
2b
3
1
(a)
Left atrium
Right atrium
Left ventricle
Right ventricle
Ventricular filling
Atrial contraction
Isovolumetric contraction phase
Ventricular ejection phase
Isovolumetric relaxation
Ventricular filling
1
2a
2b
3
1
Ventricular filling (mid-to-late diastole)
Ventricular systole (atria in diastole)
Early diastole
(b)


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Changes during Cardiac cycle


- Volume changes: End-diastolic volume, End-systolic volume, Stroke volume and Cardiac output.
- Aortic pressure: Diastolic pressure ~80 mmHg, Systolic pressure ~ 120 mmHg, most of systole ventricular pressure higher than aortic
- Ventricular pressure: Diastolic ~ 0, systolic Lt. ~120 Rt. ~ 25 mmHg.
- Atrial pressure: A wave =atrial systole, C wave= ventricular contraction (AV closure), V wave= ventricular diastole (Av opening)
- Heart sounds: S_1 = turbulence of blood around a closed AV valves, S_2 = turbulence of blood around a closed semilunar valves.

Heart Sounds



Heart Sounds

- Heart sounds (lub-dup) are associated with closing of heart valves



Heart sounds

- Auscultation – listening to heart sound via stethoscope
- Four heart sounds
  - S_1 – "lubb" caused by the closing of the AV valves
  - S_2 – "dupp" caused by the closing of the semilunar valves
  - S_3 – a faint sound associated with blood flowing into the ventricles
  - S_4 – another faint sound associated with atrial contraction

Cardiac Output (CO) and Reserve

- CO is the amount of blood pumped by each ventricle in one minute
- CO is the product of heart rate (HR) and stroke volume (SV)
- HR is the number of heart beats per minute
- SV is the amount of blood pumped out by a ventricle with each beat
- Cardiac reserve is the difference between resting and maximal CO

Cardiac Output: Example

- CO (ml/min) = HR (75 beats/min) x SV (70 ml/beat)
- CO = 5250 ml/min (5.25 L/min)

Regulation of Stroke Volume


- SV = end diastolic volume (EDV) minus end systolic volume (ESV)
- EDV = amount of blood collected in a ventricle during diastole
- ESV = amount of blood remaining in a ventricle after contraction

Factors Affecting Stroke Volume


- Preload – amount ventricles are stretched by contained blood
- Contractility – cardiac cell contractile force due to factors other than EDV
- Afterload – back pressure exerted by blood in the large arteries leaving the heart

Frank-Starling Law of the Heart

- Preload, or degree of stretch, of cardiac muscle cells before they contract is the critical factor controlling stroke volume
- Slow heartbeat and exercise increase venous return to the heart, increasing SV
- Blood loss and extremely rapid heartbeat decrease SV



(a) Preload



(b) Afterload

Cardiac Output

Increased end-diastolic volume (stretches the heart)

Positive inotropic agents such as increased sympathetic stimulation; catecholamines, glucagon, or thyroid hormones in the blood; increased Ca^{2+} in extracellular fluid

Decreased arterial blood pressure during diastole

Increased PRELOAD

Increased CONTRACTILITY

Decreased AFTERLOAD

Within limits, cardiac muscle fibers contract more forcefully with stretching (Frank–Starling law of the heart)

Positive inotropic agents increase force of contraction at all physiological levels of stretch

Semilunar valves open sooner when blood pressure in aorta and pulmonary artery is lower



Increased STROKE VOLUME

Increased CARDIAC OUTPUT

Increased HEART RATE

Increased sympathetic stimulation and decreased parasympathetic stimulation

Catecholamine or thyroid hormones in the blood; moderate increase in extracellular Ca^{2+}

Infants and senior citizens, females, low physical fitness, increased body temperature

NERVOUS SYSTEM
Cardiovascular center in medulla oblongata receives input from cerebral cortex, limbic system, proprioceptors, baroreceptors, and chemoreceptors

CHEMICALS

OTHER FACTORS

Extrinsic Factors Influencing Stroke Volume


- Contractility is the increase in contractile strength, independent of stretch and EDV
- Increase in contractility comes from:
  - Increased sympathetic stimuli
  - Certain hormones
  - Ca^{2+} and some drugs


Extrinsic Factors Influencing Stroke Volume

- Agents/factors that decrease contractility include:
  - Acidosis
  - Increased extracellular K^+
  - Calcium channel blockers

Contractility and Norepinephrine


- Sympathetic stimulation releases norepinephrine and initiates a cyclic AMP second-messenger system





Potential Energy (PE)

LEFT VENTRICULAR PRESSURE/VOLUME P/V LOOP




Valvular Function

- To prevent back-flow.
- Chordae tendineae are attached to A-V valves.
- Papillary muscle, attached to chordae tendineae, contract during systole and help prevent back-flow.
- Because of smaller opening, velocity through aortic and pulmonary valves exceed that through the A-V valves.


Valvular Function (cont'd)

- Most work is external work or pressure-volume work.
- A small amount of work is required to impart kinetic energy to the heart ($1/2\ mV^2$).
- What is stroke-volume in previous figure?
- External work is area of Pressure-Volume curve.
- Work output is affected by "preload" (end-diastolic pressure) and "afterload" (aortic pressure).


Work Output of the Heart




A


Left Ventricular Pressure
Increased
preload
3
2
1
4
Left Ventricular Volume


B


Left Ventricular Pressure
Increased
afterload
3
2
1
4
Left Ventricular Volume

C


Left Ventricular Pressure
Increased
contractility
3
2
1
4
Left Ventricular Volume

PRESSURE/VOLUME RELATIONSHIPS UNDER DIFFERENT CONDITIONS




PRELOAD

AFTERLOAD

CONTRACTILITY

Intrinsic Control of Stroke Volume (Frank-Starling Curve)



Regulation of Heart Rate

- Positive chronotropic factors increase heart rate
- Negative chronotropic factors decrease heart rate


Regulation of Heart Rate: Autonomic Nervous System

- Sympathetic nervous system (SNS) stimulation is activated by stress, anxiety, excitement, or exercise
- Parasympathetic nervous system (PNS) stimulation is mediated by acetylcholine and opposes the SNS
- PNS dominates the autonomic stimulation, slowing heart rate and causing vagal tone

Chemical Regulation of the Heart

- The hormones epinephrine and thyroxine increase heart rate
- Intra- and extracellular ion concentrations must be maintained for normal heart function

CARDIAC OUTPUT CURVES



Ventricular Pressure and Volume Curves (cont'd)

- During the latter part of the ejection phase how can blood still leave the ventricle if pressure is higher in the aorta? Momentum of blood flow
- Total energy of blood = $P + mV^2/2$

  = pressure + kinetic energy
- Total energy of blood leaving ventricle is greater than in aorta.

Ejection Fraction

- End diastolic volume = 125 ml
- End systolic volume = 55 ml
- Ejection volume (stroke volume) = 70 ml
- Ejection fraction = 70ml/125ml = 56%
  (normally 60%)
- If heart rate (HR) is 70 beats/minute, what is cardiac output?
- Cardiac output = HR * stroke volume
  = 70/min. * 70 ml
  = 4900ml/min.

Ejection Fraction (cont'd)

- If HR =100, end diastolic volume = 180 ml, end systolic vol. = 20 ml, what is cardiac output?
- C.O. = 100/min. * 160 ml = 16,000 ml/min.
- Ejection fraction= 160/180%=~ 90%


Aortic Pressure Curve


- Aortic pressure starts increasing during systole after the aortic valve opens.
- Aortic pressure decreases toward the end of the ejection phase.
- After the aortic valve closes, an *incisura* occurs because of sudden cessation of back-flow toward left ventricle.
- Aortic pressure decreases slowly during diastole because of the elasticity of the aorta.

Frank-Starling Mechanism

- Within physiological limits the heart pumps all the blood that comes to it without excessive damming in the veins.
- Extra stretch on cardiac myocytes makes actin and myosin filaments interdigitate to a more optimal degree for force generation.

Ventricular Stroke Work Output





Autonomic Effects on Heart


- Sympathetic stimulation causes increased HR and increased contractility with HR = 180-200 and C.O. = 15-20 L/min.
- Parasympathetic stimulation decreases HR markedly and decreases cardiac contractility slightly. Vagal fibers go mainly to atria.
- Fast heart rate (tachycardia) can decrease C.O. because there is not enough time for heart to fill during diastole.

At HR 72 bpm, systole occupies 40% of the cardiac cycle. At HR 200 bpm, it occupies 65% of the time.

Each person has his own optimal maxHR.

HR	Systole	Diastole
60	0.3	0.7
75	0.25	0.55
200	0.19	0.11

Effect of Sympathetic and Parasympathetic Stimulation on Cardiac Output



Thank You

